

Antimicrobial Activity of Honey with Special Reference to Methicillin Resistant *Staphylococcus aureus* (MRSA) and Methicillin Sensitive *Staphylococcus aureus* (MSSA)

GURUVU NEERAJA RANI¹, RADHIKA BUDUMURU², NARASINGA RAO BANDARU³

ABSTRACT

Introduction: Antimicrobial agents are essentially important in reducing the global burden of infectious diseases. With the irrational and excessive use of antibiotics in underdeveloped and developing countries, there may be chances to develop and spread resistant pathogens in the community. As a result, the effectiveness of the antibiotics is diminishing. Therefore, the need for novel alternative antimicrobial strategies has renewed interest in natural products like turmeric, honey, ginger and others exhibiting antibacterial properties. This situation has led to a re-evaluation of the therapeutic use of ancient remedies like honey as no other studies are available in the state of Andhra Pradesh with the locally available honey.

Aim: To find out the efficacy of antibacterial activity of locally available honey against Methicillin Resistant *Staphylococcus aureus* (MRSA) and Methicillin Sensitive *Staphylococcus aureus* (MSSA) isolates.

Materials and Methods: A prospective study on the antibacterial activity of Bharat multi floral pasteurised honey which was locally available in the state of Andhra Pradesh, further it was conducted and evaluated against the bacterial strains of Methicillin Resistant *Staphylococcus aureus* and Methicillin Sensitive *Staphylococcus aureus*. Their antibacterial sensitivity pattern was tested using Kirby-Bauer disc diffusion susceptibility testing technique of CLSI along with other commonly used antimicrobials.

Results: Both MRSA and MSSA isolates were sensitive to honey. But MRSA were resistant to all antimicrobials tested except linezolid where as MSSA were sensitive to all except penicillin.


Conclusion: It is definitely worthy to consider honey as a promising future antimicrobial to be tested and studied. Honey, may be elaborately used in future with some more molecular studies on its method of action as an antimicrobial agent.

Keywords: Antibacterial activity, Kirby-bauer disc diffusion, Minimum inhibitory concentration, Zone of Inhibition


INTRODUCTION

Antimicrobial agents are essentially important in reducing the global burden of infectious diseases [1]. The selection and spread of multi-resistant organisms in developing countries, which can often be traced to complex socio-economic and behavioural factors, contribute to the escalating problem of antibiotic resistance worldwide. In developing countries, the irrational use of antibiotics by health professionals, unskilled practitioners, laypersons, poor drug quality, unhygienic conditions and inadequate surveillance account for the spread of resistant bacteria. Nevertheless, misuse of antibiotics is one of the reasons for the increasing rates of resistance, especially in rural areas [2]. There is an emergence of drug resistance for the drugs like vancomycin and daptomycin leading to search for still newer drugs for combating the drug resistance in Staphylococci [3]. With the irrational and excessive use of antibiotics in underdeveloped and developing countries the developed resistance may spread in the community making the strains as super bugs causing difficulties in eradication [4]. As a result, the effectiveness of the antibiotics is diminished [5]. Therefore, the need for novel alternative antimicrobial strategies has renewed interest in natural products like turmeric, honey, ginger etc., exhibiting antibacterial properties. This situation has led to a re-evaluation of the therapeutic use of ancient remedies including honey [6-8]. Honey is well known as a magic drug for almost all kinds of diseases, not to mention the fact that many people do depend more on folk medicine and natural remedies which are cheap that have been known for their therapeutic effects over the past decades [6]. Honey has well established function as

an effective antibacterial agent with a broad spectrum of activity against Gram-positive and Gram-negative bacteria [9-11]. The application of honey can promote the healing of infected wounds that do not respond to the conventional therapy, i.e., antibiotics and antiseptics [12] including wounds infected with methicillin-resistant *S. aureus* [13,14]. Laboratory studies have revealed that honey is effective against MRSA, β -haemolytic streptococci and Vancomycin Resistant Enterococci (VRE) [15,16]. The beneficial role of honey is attributed to its antibacterial property with regards to its high osmolarity, acidity (low pH) and content of hydrogen peroxide (H_2O_2) and non-peroxide components, i.e., the presence of phytochemical components like Methylglyoxal (MGO). The antimicrobial agents in honey are predominantly hydrogen peroxide, of which the



[Table/Fig-1]: Antimicrobial activity of honey and other common antibacterial agents on/against MRSA.



[Table/Fig-2]: Antimicrobial activity of honey and other common antibacterial agents on/against MSSA.

H=Honey, AMC = Amoxicillin + Clavulanic Acid, CTR = Ceftriaxone, VA = Vancomycin, P = Penicillin, LZ = Linezolid, GEN = Gentamicin, Cefperazone (CPZ), PTZ = Piperacillin + Tazobactam, CAC = Ceftazidime + Clavulanic Acid, CZ = Cefazolin, CX = Cefoxitine

Zone of Inhibition of Honey on/against MRSA = 36.2± 0.2

Zone of Inhibition of Honey on/against MSSA = 40.16±0.152

concentration is determined by relative levels of glucose oxidase, synthesized by the bee and catalase originating from flower pollen [17]. Hence, an attempt was made to find out the efficacy of locally available honey against MRSA which emerged as a super bug and MSSA and their antibacterial activity to commonly used antimicrobials.

MATERIALS AND METHODS

The antibacterial activity of Bharat multi floral pasteurized honey obtained from Bharat Unani Pharmacy (Bharat honey co), Hyderabad, Andhra Pradesh, India, was tested and evaluated in the month of April 2017 against the bacterial strains of MRSA and MSSA obtained from Saveetha University, Chennai, India [3] which were already confirmed phenotypically (Disc diffusion test with Cefoxitin 30 µg disc (obtained by Himedia, Mumbai) was used to differentiate MRSA and MSSA isolates and the interpretation was done by MSSA if zone size was β 22 mm. The strain was considered as MRSA if zone size was <22 mm. (ref. CLSI M100-S23) [18] and genotypically (DNA sequencing was done for MecA gene at Eurofins Genomics India Pvt., Ltd. Bangalore, Karnataka, India). Their antibacterial sensitivity pattern was tested using Kirby-Bauer disc diffusion susceptibility testing technique of CLSI [18]. MRSA ATCC strain No. 43300, MSSA ATCC 25923 were included as a positive control strains and MRSA ATCC 33591, MSSA ATCC 29213 were used as negative control strains. A six hour incubated bacterial culture suspension matching with 0.5 Mc-Farland scale standard was prepared equivalent to 1.5×10^8 CFU/ ml organisms in 5 ml peptone water and spread onto the sterile Mueller-Hinton agar (Himedia, Mumbai) plates to prepare a lawn culture. Dried in the incubator for half an hour and three such plates were prepared for

each bacterial strain. Honey disks were prepared by using Watman No.1 filter paper and discs were punched with a office paper hole punching machine of 6 mm diameter and such 100 discs were taken in small glass bottle and sterilized at 160°C for two hours in a hot air oven. To these 100 sterile discs, 1 ml of pasteurized 100% V/V undiluted honey obtained for this study was added and kept for overnight for equal absorption of honey by all discs [19]. Standard antibiotic discs of Amoxicillin+ Clavulanic Acid (AMC) (20/10 µg), Ceftriaxone (CTR) (30 µg), Vancomycin (VA) (30 µg), Penicillin (P) (10 Units), Linezolid (LZ) (30 µg), Gentamicin (GEN) (10µg), Cefoperazone (CPZ) (30 µg), Piperacillin+tazobactam (PTZ) (100/10 µg), Cefpazidime+clavulanicacid (CAC) (30/10 µg), Cefazolin (CZ) (30 µg) and Cefoxitine (CX) (30 µg) obtained from Himedia laboratories Pvt Ltd., Mumbai, India, and the prepared honey discs were placed aseptically on the Mueller Hinton agar. Plates were left for one hour at 25°C to allow a period of preincubation diffusion in order to minimize the effect of variation in time between the placements of different discs. The plates were then incubated aerobically at 37°C over night to allow bacterial growth. After incubated plates were observed and the zone of inhibition was measured to evaluate the antimicrobial activity for each of the tested antibiotics and honey samples using a special scale obtained from Himedia Laboratories, Mumbai, India. The sensitivity testing plates were done in triplicates for each strain of MRSA and MSSA isolates and the zone of inhibition were measured to the nearest millimeters. To calculate the mean and standard deviation of each strain statistically using Statistical Package for the Social Sciences (SPSS) software. We took the criteria for sensitivity was as the zone of inhibition for honey is bigger than the zones any of the antimicrobials used for testing the MRSA and MSSA isolates.

The present study was approved by the Institutional Ethical Committee.

RESULTS

Both MRSA and MSSA were sensitive to honey with a zone of inhibition of 36.2±0.2 mm and 40.16±0.152 mm respectively. But MRSA were resistant to all antimicrobials tested except linezolid (21.03±0.152 mm) whereas, MSSA were sensitive to all except penicillin with no zone of inhibition [Table/Fig-1-3].

DISCUSSION

It has been reported that honey showed both bacteriostatic and bactericidal effect against many Gram-positive as well as Gram-negative bacteria [20-24]. The use of natural products to enhance wound healing is a common practice in many parts of the world. Honey consists of a super saturated solution of sugars and has a low pH between 3.2 and 4.5. This pH together with honey's high

Name of Organism	Diameter of Zone of Inhibition (mm)											
	Honey (H)	Amoxicillin + Clavulanic Acid (AMC) (20/10 µg)	Ceftriaxone (CTR) (30 µg)	Vancomycin (VA) (30 µg)	Penicillin (P) (10 Units)	Linezolid (LZ) (30 µg)	Gentamicin (GEN) (10µg)	Cefoperazone (CPZ) (30 µg)	Piperacillin + Tazobactam (PTZ) (100/10 µg)	Ceftazidime + Clavulanic Acid (CAC) (30/10 µg)	Cefazolin (CZ) (30 µg)	Cefoxitine (CX) (30 µg)
Methicillin resistant <i>Staphylococcus aureus</i> (MRSA)	36.2± 0.2	8.2± 0.2	16.13± 0.152	10.2± 0.2	0	21.03± 0.152	22.13± 0.152	10.1± 0.1	12± 0.1	8.166± 0.152	18.16± 0.152	0
	(S)	(R)	(R)	(R)	(R)	(S)	(R)	(R)	(R)	(R)	(R)	(R)
Methicillin sensitive <i>Staphylococcus aureus</i> (MSSA)	40.16± 0.152	23.03± 0.152	40.1± 0.1	23.03± 0.15	0	45.13± 0.152	44.1± 0.1	44.13± 0.152	46.16± 0.152	30± 0.1	48.16± 0.152	27.96± 0.152
	(S)	(S)	(S)	(S)	(R)	(S)	(S)	(S)	(S)	(S)	(S)	(S)

[Table/Fig-3]: Antimicrobial sensitivity pattern of MRSA and MSSA with reference to honey and other common antimicrobials.

H=Honey, AMC = Amoxicillin + Clavulanic Acid, CTR = Ceftriaxone, VA = Vancomycin, P = Penicillin, LZ = Linezolid, GEN = Gentamicin, Cefperazone (CPZ), PTZ = Piperacillin + Tazobactam, CAC = Ceftazidime + Clavulanic Acid, CZ = Cefazolin, CX = Cefoxitine

Author/s	Ref.No.	Type of honey/Concentration	MRSA-Sensitivity/Zone of Inhibition	MSSA- Sensitivity/Zone of Inhibition
Patel A et al.,	[4]	Karapur village honey from Goa	Sensitive/NA	Sensitive/NA
Almasaudi SB et al.,	[34]	Manuka /20%,30%,40% V/V Nigella Sativa/20%,30%,40% V/V Sidr/20%,30%,40% V/V	Sensitive/NA Sensitive/NA Sensitive/NA	Sensitive/NA Sensitive/NA Sensitive/NA
Zakaria AS	[35]	Yemen Sidr Honey (YSH)/100% V/V Southern Sidr Honey (SSH) 100% V/V Multi-Flower Mountain Honey (MMH) 100% V/V	Sensitive/14±2.83 Sensitive/15±0.71 Sensitive/9±1.1	Sensitive/15±2.83 Sensitive/17±1.35 Sensitive/13±3.24
Neeraja Rani G et al.,	Present Study	Bharat Honey from Andhra Pradesh / 100% V/V	Sensitive/36.2± 0.2	Sensitive/40.16±0.152

[Table/Fig-4]: Comparative study of Antimicrobial activity of honey on/against MRSA and MSSA.

NA = Not available

osmolarity and the presence of H_2O_2 reduces the bacterial growth at the wound site. Honey in wound dressing has been reported to provide ideal environment for the rapid tissue repair and regeneration that are essential for growth of wound bed [25]. *Staphylococcus aureus* is the most frequently isolated wound pathogen and it is becoming increasingly resistant to antibiotics in common use. Honey has been reported to be effective in eradicating antibiotic resistant bacteria including MRSA [26] which is a super bug now. Any zone diameter having less than 7 mm shows that the organism is resistant to the honey sample but if the zone diameter is greater than 11 mm it suggests that the microorganism is sensitive to the honey with special reference to *Pseudomonas aeruginosa* [27]. The findings of our study together with four other previous studies [28-30] show that honey promises to be an effective wound antiseptic with broad spectrum antimicrobial activity. Some topical antimicrobials adversely affect the human skin/tissue and repair process during the treatment of wounds where as there is no need for laboratory evaluation of honey as it does not adversely affect human skin/tissue [31]. The special character of honey is the potential to limit the growth of wound pathogens, but also there is evidence that honey has the potential to promote the healing [32,33] and no other antimicrobial agent possesses these characteristics. Honey is effective even it is diluted by burn wound exudates. In burns, honey's antimicrobial and anti inflammatory properties allow a moist healing environment to be maintained that protects the wounds from deterioration and fibrosis [29]. [Table/Fig-1-3] shows the results of antibacterial activity of honey towards the two microorganisms tested. MRSA as well as MSSA were sensitive to undiluted honey samples tested with an average zone of inhibition of 36.2+0.2 and 40.16+ 0.152 mm respectively. Ogbaje EO et al., and Murthy K et al., found in their study that 100% V/V undiluted honey inhibited the growth of MRSA with a zone of inhibition of 18 mm and 11 mm respectively [22,33], whereas Patel A et al., in their study observed that diluted honey of 20% V/V, 30% V/V and 40% V/V inhibited the growth of MRSA [4] whereas, 15% V/V also inhibited the growth of MSSA in addition to other concentrations used for MRSA and Almasaudi SB et al., observed the only 50% V/V concentrated honey inhibited the growth of both MRSA and MSSA [34], whereas, Zakaria AS showed that 100% V/V undiluted honey inhibited MRSA with a zone of 14±2.83 mm and MSSA 15±2.83 mm for Yemen Sidr honey [35], MRSA with a zone of 15±0.71 mm and MSSA 17±1.35 mm for Southern Sidr honey and MRSA with a zone of 9±1.1 mm and MSSA 13±3.24 mm for multi-flower mountain honey. In our findings with our local honey, the zone of inhibitions of MRSA (36.2+0.2) and MSSA (40.16+0.152) were much bigger than the ones reported by other workers. Hence, we can say that the efficacy of our local honey was more than the honey samples of other workers. Neerajarani G et al., [36] in their study showed the antibacterial activity of undiluted honey on the isolates of *Staphylococcus aureus*, *E. coli* and *Pseudomonas aeruginosa* and all these organisms were sensitive to undiluted honey. The exact explanation for the antibacterial activity of honey is not known, but it is clear that the higher the concentration of honey the greater its usefulness as an antibacterial agent. However, it is expected that

the clinical significance of the antibacterial activity in honey will be unequivocally proven only if a clinical trial is conducted to compare dressings of different sugars and selected honeys. Although more research is needed, as with many of the therapeutic interventions used in modern wound care, in the absence of data from well controlled clinical trials. Recent reviews on the successful usage of honey as a dressing on infected wounds show that many authors support the use of honey in infected wounds and some suggest the prophylactic use of honey on the wounds of patients susceptible to MRSA and other antibiotic-resistant bacteria [37]. Well documented clinical trials and researches are going on honey and nanotechnology which may provide promising results on therapeutic use of honey in the future. Comparative studies of antimicrobial activity of honey on/against MRSA and MSSA were shown in [Table/Fig-4]. Further in continuation of studies on honey we are planning to study the efficacy of honey with nano particles of gold, silver on the wound healing process in an experimental animal rat.

LIMITATION

The present study was limited to testing of 100% V/V of local honey. Testing of different dilutions is under study.

CONCLUSION

It is definitely worthy to consider honey as a promising future antimicrobial to be tested and studied. Rediscovering honey as a natural remedy for wound pathogens proved its effectiveness on antimicrobial resistant strains of bacteria including MRSA. In the present study, we tried to focus more on whether honey can be used for treating Staphylococcal infections with special reference to MRSA. Honey, the nature blessed and environmental friendly product may be elaborately used in future with some more molecular studies on its method of action as an antimicrobial agent.

REFERENCES

- [1] Deb Mandal M, Mandal S. Honey: its medicinal property and antibacterial activity. *Asian Pac J Trop Biomed*. 2011;1(2):154-60.
- [2] Okeke IN, Lamikanra A, Edelman R. Socioeconomic and behavioral factors leading to acquired bacterial resistance to antibiotics in developing countries. *Emerging Infectious Diseases*. 1999;5(1):18-27.
- [3] Srinivas B, Murthy SN, Prasad U, Rao BN. Demographic profile and drug resistance pattern in methicillin sensitive staphylococcus aureus in rural and urban tertiary care centers. *Journal of Pure and Applied Microbiology*. 2015;9(4):2961-69.
- [4] Patel A, Chauhan BP. Antimicrobial effect of Honey on MRSA isolated from pus samples. *Int J Drug Res Tech*. 2016;6(2):58-63.
- [5] Levy SB, Marshall B. Antibacterial resistance worldwide: causes, challenges and responses. *Nat Med*. 2004;10:122-29.
- [6] Bagde AB, Sawant RS, Bingare SD, Sawai RV, Nikumbh MB. Therapeutic and nutritional values of honey [Madhu]. *Int Res J Pharm*. 2013;4(3):19-22.
- [7] Mandal S, Deb Mandal M, Pal NK. Synergistic anti- *Staphylococcus aureus* activity of amoxicillin in combination with *Embilica officinalis* and *Nymphaea odorata* extracts. *Asian Pac J Trop Med*. 2010;3(9):711-14.
- [8] Mandal S, Deb Mandal M, Pal NK, Saha K. Antibacterial activity of honey against clinical isolates of *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella enterica* serovar Typhi. *Asian Pac J Trop Med*. 2010;3(12):961-64.
- [9] Irish J, Blair S, Carter DA. The antibacterial activity of honey derived from Australian flora. *PLoS ONE*. 2011;6:e18229.
- [10] Molan PC. The antibacterial nature of honey. The nature of the antibacterial activity. *Bee World*. 1992;73(1):5-28.

- [11] Tan HT, Rahman RA, Gan SH, Halim AS, Hassan SA, Sulaiman SA, et al. The antibacterial properties of Malaysian tualang honey against wound and enteric microorganisms in comparison to manuka honey. *BMC Complement Alternat Med.* 2009;9:34.
- [12] Ahmed AK, Hoekstra MJ, Hage J, Karim RB. Honey-medicated dressing: transformation of an ancient remedy into modern therapy. *Ann Plast Surg.* 2003;50:143–48.
- [13] Natarajan S, Williamson D, Grey J, Harding KG, Cooper RA. Healing of an MRSA-colonized hydroxyurea-induced leg ulcer with honey. *J Dermatol Treat.* 2001;12:33–36.
- [14] Dunford C, Cooper RA, Molan PC. Using honey as a dressing for infected skin lesions. *Nurs Times.* 2000;96:7–9.
- [15] Allen KL, Hutchinson G, Molan PC. Potential for using honey to treat wounds infected with MRSA and VRE. *First World Healing Congress, Melbourne, Australia.* 2000:10–13.
- [16] Kingsley A. The use of honey in the treatment of infected wound. *British J Nursing.* 2001;10:13–16.
- [17] Weston RJ. The contribution of catalase and other natural products to the antibacterial activity of honey: a review. *Food Chemistry.* 2000;71:235–39.
- [18] CLSI. 2013. Performance standards for antimicrobial susceptibility testing. CLSI approved standard M100-S23. Clinical and Laboratory Standards Institute, Wayne, PA.
- [19] Cheesbrough M. District laboratory practice in tropical countries. Part II. Low price edition Biochemical test to identify bacteria; antimicrobial susceptibility testing. The Cambridge University Press; 2000.
- [20] Badawy OFH, Shafii SSA, Tharwat EE, Kamal AM. Antibacterial activity of bee honey and its therapeutic usefulness against *Escherichia coli* O157:H7 and *Salmonella typhimurium* infection. *Rev Sci Tech Off Int Epiz.* 2004;23(3):1011–22.
- [21] Wilkinson JM. Antibacterial activity of 13 honeys against *Escherichia coli* and *Pseudomonas aeruginosa*. *J Med Food.* 2005;8:100–03.
- [22] Ogbaje EO, Ogansanya T, Aiwerioba OIR. Conventional use of honey as antibacterial agent. *Ann Afri Med.* 2006;5(2):78–81.
- [23] Agbagwa OE, Peterside NF. Effect of raw commercial honeys from Nigeria on selected pathogenic bacteria. *Afr J Microbiol Res.* 2010;4(16):1801–03.
- [24] Othman AS. Antibacterial activity of bee and Yemeni Sidr honey against some pathogenic bacterial species. *Int J Curr Microbiol App Sci.* 2014;3(10):1015–25.
- [25] Molan P, Rhodes T. Honey: a biological wound dressing. *Wounds* 2015;27(6):141–51.
- [26] Nagi AA, Amghalia E, Shamsudin MN, Abdullah R, Mohammed R, Sekawi Z. Antibacterial activity of honey against methicillin resistant *Staphylococcus aureus*. *Research journal of Biological Sciences.* 2009;4(8):943–47.
- [27] Cooper RA, Molan PC. The use of honey as an antiseptic in managing *Pseudomonas* infection. *J Wound Care.* 1999;8:161–64.
- [28] Cooper RA, Molan PC, Harding KG. Antibacterial activity of honey against strains of *Staphylococcus aureus* from infected wounds. *J of Royal Soc Med.* 1999;92:283–85.
- [29] Cooper RA, Molan PC, Harding KG. The sensitivity to honey of Gram-positive cocci of clinical significance isolated from wounds. *J Applied Microbiol.* 2002;93:857–63.
- [30] Ward RS, Saffle JR. Topical agents in burn and wound care. *Phys Ther.* 1995;75:526–38.
- [31] Molan PC. The role of honey in the management of wounds. *J Wound Care.* 1999;8:415–18.
- [32] Tonks A, Cooper RA, Price AJ, Molan PC, Jones KP. Stimulation of TNF- α release in monocytes by honey. *Cytokine.* 2001;14:240–42.
- [33] Murthy K, Ramya SR, Roplekar P, Shah D, Sharma N. To study the antibiotic susceptibility of the isolated strains of *Staphylococcus aureus* and comparative analysis with natural herbs. *International Science Journal.* 2014;1(3):13–18.
- [34] Almasaudi SB, Al-Nahari AAM, El-Ghany ESMA, Barbour E, Al-Muhayawi SM, Al-Jaouni S, et al. Antimicrobial effect of different types of honey on *Staphylococcus aureus*. *Saudi Journal of Biological Sciences* 2016 in press. Available at: <http://www.sciencedirect.com/science/article/pii/S1319562X16300870>. Last accessed on 18.07.2017.
- [35] Zakaria AS. Mechanism of antibacterial action of honey on pathogenic wound bacterial strains: A proteomic analysis. *Int Res J Pharm.* 2015;6(11):778–88.
- [36] Neerajarani G, Rao BN, Sukumar E, Padmaja J, Misra AK. A prospective study of honey with special reference to its antibacterial activity. *International Journal of General Medicine and Pharmacy.* 2016;5(5):60–73.
- [37] Molan PC. Honey as a topical antibacterial agent for treatment of infected wounds. *World Wide Wounds.* 2011. Available at: <http://www.worldwidewounds.com/2001/november/Molan/honey-as-topical-agent.html>. Last accessed on 18.07.2017.

PARTICULARS OF CONTRIBUTORS:

1. Research Scholar, Saveetha University, Chennai, Tamil Nadu, India.
2. Assistant Professor, Department of Microbiology, Great Eastern Medical School, Srikakulam, Andhra Pradesh, India.
3. Professor and Head, Department of Microbiology, Gayatri Vidyaparishad Institute of Health Care and Medical Technology, Visakhapatnam, Andhra Pradesh, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Narasinga Rao Bandaru,
16-4-21, Sri Sai Sadan, Official Colony, Maharani-peta, Visakhapatnam-530002, Andhra Pradesh, India.
E-mail: narasinga_bandaru@rediffmail.com

Date of Submission: **May 15, 2017**

Date of Peer Review: **Jun 07, 2017**

Date of Acceptance: **Jul 20, 2017**

Date of Publishing: **Aug 01, 2017**

FINANCIAL OR OTHER COMPETING INTERESTS: None.