

Randomized Controlled Trial Comparing Open, Conventional, and Single Port Laparoscopic Appendectomy

VARUN KUMAR SINGH¹, KUMAR NISHANT², BIKRAM KHARGA³,
ABONI KUMAR KALITA⁴, PHUCHUNGLA BHUTIA⁵, JATIN JAIN⁶

ABSTRACT

Introduction: Open appendectomy took decades to evolve to establish itself as the treatment for appendicitis. Then came the era of laparoscopy, which unlike in case of cholecystectomy, faced quite resistance and still has not become the treatment of choice. What followed was the reduction in the size and number of ports. Single port laparoscopic appendectomy and Natural Orifice Transluminal Endoscopic Surgery (NOTES) were the further developments.

Aim: To evaluate the status of appendectomy in patients with simple appendicitis through the three approaches: Open Appendectomy i.e. OA, Conventional Laparoscopy Appendectomy i.e. CLA and Single Port Laparoscopic Appendectomy (SPLA)/Incision Approach.

Materials and Methods: The study was designed as a double blinded randomised controlled trial and included the patients aged 12-50 years who had simple uncomplicated appendicitis and underwent appendectomy over a duration of 18 months.

Results: Mean age of patients was 31.27, 27.4 and 27.64 years and female to male ratio was 1.14, 1.5 and 1.33 for OA, CLA and SPLA arm, respectively. Overall duration of presentation was two days and most of the patients were under BMI of 25. The appendix was the first visualized organ in less than half of

the total patients and a little over half had adhesions. The most common location of the appendix was paracaecal followed by pelvic. There were very few intraoperative complications like bleeding from appendicular artery and spillage from appendix in one patient who underwent OA. Mean duration of surgery was 65.18 minutes (SPLA>CLA>OA). Surgery took less time in patients with BMI<25 (SPLA>CLA>OA). CLA took substantially less time (49.5 minutes) in patients with BMI≥25 (CLA>OA>SPLA). Pain was significantly higher in minimal invasive procedures compared to OA in immediate postoperative period, which settled 4th hour onwards and remained on higher side for patients who underwent OA. Three fourth patients were able to accept orally after six hours. Wound infection rate was low (OA>CLA=SPLA). Patients who underwent OA stayed in hospital for a longer time and resumed their duty much later compared to minimal invasive arm. Scar assessment score and overall satisfaction were not much different among the patients with simple appendicitis of three arms. Cost of treatment was significantly higher for SPLA compared to other treatment arms.

Conclusion: In patients with simple uncomplicated appendicitis, OA, CLA and SPLA do not differ much in outcome especially in lean and thin patients.

Keywords: Appendicitis, Cholecystectomy, Natural orifice transluminal endoscopic surgery

INTRODUCTION

Since accurate etiologies remained elusive over the centuries, treatment also continued to remain so. First appendectomy was done by Amyand in 1735, when he was operating for an inguinal hernia [1]. It took another 150 years before early appendectomy was advocated as a treatment for this condition. Various modifications were proposed, accepted and rejected over next several decades, but most noticeable was the advent of laparoscopic approach. Yet, laparoscopic approach was never as well embraced by surgeons as its cholecystectomy counterpart. As patients transcended into clients, further developments were sought, initial step was improving patient satisfaction by reducing size and number of ports. With technological improvements in laparoscopic imaging equipment and instruments this need was first fulfilled by a laparoscopic appendectomy through a single umbilical incision. Progression to 'scarless' surgery through natural orifices is the next logical step. Several studies have compared open and laparoscopic

cholecystectomy, while others have compared different laparoscopic approaches. However, there seems to a paucity of literature comprehensively comparing open, conventional laparoscopic and single incision laparoscopic appendectomy in a prospectively conducted RCT providing an evidence of satisfactory quality. This study was done in an attempt to comprehensively address these three approaches to appendectomy, and encourage adoption of laparoscopic approach in this remote part of the country, if found to be superior.

MATERIALS AND METHODS

The study was designed as a prospective double blinded randomized controlled comparison of three treatment modalities [designated as intervention groups] viz OA, CLA and SPLA. The study was conducted in Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim in North East India for a duration of 18 months from January 2014 to June 2015. The study was approved by the

research protocol evaluation committee and ethical clearance was obtained from the Institutional Ethical Committee.

All the patients who underwent appendectomy formed the population group. Inclusion criteria into the study group were patients willingness for randomisation, diagnosis of 'simple' appendicitis supported by ultrasonography and age between 12 and 50 years. Any contraindication to general anaesthesia, comorbidities other than hypertension and diabetes mellitus, 'complicated' appendicitis [perforation, abscess, and lump] and technical glitches leading to ergonomic difficulties were major exclusions.

An audio visual interactive session of 15 minutes about appendicitis, appendectomy and process of randomization was planned with the patients and their relatives to convince them and help them reaching a decision. They were communicated that they could choose procedure of their choice but if consented to be included in the study they would be randomized and might not know about what procedure they had undergone until 24 hours following surgery.

All patients were operated under general anaesthesia and abdomen was covered with a large dressing which was removed only after 24 hours to ensure blinding at patients end. They were also assured that they would get the standard care as per hospital protocol irrespective of their decision for inclusion into the study. Patients were then randomized using computer generated random numbers in to three intervention arms. Blinding was also ensured at investigators end by forming four teams, 1st for counselling and randomization, 2nd for operative intervention, 3rd for recording of data and last for analysis. All three operative interventions were standardized by a

team of experts in the department of surgery after careful review of literature. Operative team remained strictly adhered to protocols and sequence of steps during entire study duration. Final outcomes were shared only after the last team analyzed whole data [Table/ Fig-1 and 2].


RESULTS

Over a duration of 18 months, 186 appendectomies were performed in our institute. From this population group, 45 patients were selected to form the study group who fulfilled inclusion criteria. Patients were then randomized to undergo one of the three procedures. All treatment arms received equal number of patients but one patient in SPLA group was converted to CLA and excluded from final analysis.


There was no difference in patients characteristics when population, study group and intervention groups were matched for age, sex and BMI [Table/Fig-3].

Duration of pain or presentation, history of previous surgery, initial visualized organ on putting laparoscope, adhesions and locations of appendix were studied and have been shown in [Table/Fig-4].

Overall mean duration of surgery was a little more than an hour [65.18 minutes]. SPLA took maximum time [82.79 minutes] but difference between OA and CLA was not even one second [OA= 56.93, CLA= 57.00 minutes] [p=0.018]. Minimum duration of surgery was almost similar in all arms, least in CLA [25 minutes in CLA, 30 minutes each in OA and SPLA]. In patients with BMI <25 [n=28, 63.6%], OA took significantly less time [54 minutes] compared to CLA [62 minutes] and SPLA [89.44 minutes] [p=0.013]. In contrast, in patients with BMI >25 [n=16, 36.4%], duration of surgery was least in CLA [49.5 minutes] followed by OA and SPLA [62.8 and 70.8 minutes] [p=0.525]. Duration of surgery in patients with pain <3 days [n=33, 75%] was 61.15 minutes and in those with ≥3 days [n=11, 25%] was 77.27 minutes [p=0.110]. When treatment arms were compared within those who presented with a shorter duration pain



[Table/Fig-1]: Consort flow diagram.



[Table/Fig-2]: Flowchart showing the scheme of study.

Patient Characteristics	N		Gender*		Procedure*			
	Population	Study	Female	Male	OA	CLA	SPLA	
Age	12-20	16	2	2	0	1	1	
	21-30	74	26	16	10	9	7	
	31-40	38	15	7	8	5	4	
	41-50	22	1	0	1	1	0	
	Total	150**	44	25	19	15	15	14
	df/p [χ ²]	3/ 0.064		3/ 0.296		6/ 0.693		
	Mean	30.79	28.80	27.96	29.89	31.27	27.40	27.64
Gender	df/F/p [ANOVA]	1,192/ 1.597/ 0.205		1,42/ 1.181/ 0.283		2,41/ 2.136/ 0.131		
	Females	106	25	--	--	8	9	8
	Males	80	19	--	--	7	6	6
	Total	186	44	--	--	15	15	14
	F:M Ratio	1.32	1.31	--	--	1.14	1.5	1.33
	df/p [χ ²]	1/ 0.930		--		2/ 0.934		
	Mean	23.7	24.3	23.68	25.10	23.76	24.72	24.42
BMI	df/F/p [ANOVA]	1,228/ 1.973/ 0.161		2,41/ 1.176/ 0.284		2,41/ 0.188/ 0.829		
	<18.5	29	5	4	1	2	1	2
	18.5-25	65	23	13	10	8	8	7
	>25-30	58	11	5	6	5	4	2
	>30	34	5	3	2	0	2	3
	Total	186	44	25	19	15	15	14
	df/p [χ ²]	3/ 0.842		3/ 0.638		6/ 0.603		

[Table/Fig-3]: Patients were matched for their age, sex and BMI.


*Within the study group;

**Patients below 12 years and above 50 years have not been shown.

Characteristics/ Parameters		N	Procedure		
			OA	CLA	SPLA
Mean duration of pain [days] [p=0.609]		-	1.7	2.07	2.0
Comorbidities	Hypertension	4	2	1	1
	Diabetes	1	1	0	0
Previous operative scar ^A		2	0	1	1
Organ first visualized ^B [p=0.842]	Appendix	21	7	9	5
	Caecum	10	4	3	3
	Omentum	5	2	1	2
	Small bowel	8	2	2	4
Patients with adhesions ^C		23	8	8	7
Organ adhered to appendix ^D [p=0.738] ^E	Omentum	13	4	4	5
	Small bowel	16	7	6	3
	Anterior abdominal wall ^F	4	1	2	1
	Transverse colon	1	0	1	0
Location of appendix ^G [p=0.202]	Retrocaecal	4	1	3	0
	Paracaecal ^H	21	6	8	7
	Preileal	2	2	0	0
	Pelvic ^I	17	6	4	7
Intraoperative complications	Bleeding	1 ^J	1	0	0
	Spillage from appendix	1 ^J	1	0	0

[Table/Fig-4]: Clinical Characteristics.

^ABoth female patients, with Pfannenstiel scar; ^BChances of appendix visualization less with longer pain history [p<0.001], no impact by Gender [p=0.263] and BMI [p=0.932]; ^CPatients with BMI<25 [p=0.039] and pain ≥3 days [p=0.017] developed adhesions more commonly; ^DNo adhesions with sigmoid colon and right ovary, so excluded from final analysis; ^EPresence versus absence of adhesions; ^FMore common with pain of ≥3 days [p=0.001]; ^GNo impact on duration of pain [p=0.807]; Common in females^H and males; ^JBoth complications in same patient.



[Table/Fig-5]: Mean Visual Analog Scale Score [VASS] in postoperative period across three treatment arms. Dotted red denotes overall mean, yellow OA, blue CLA and green SPLA. Numerals against VAS indicates postoperative hour, DIS=discharge, D10=10th follow up day.

Parameter	Mean			
	OA	CLA	SPLA	p
Bowel sounds	6.27	4.8	3.86	0.336
Hospital stay	5.27	2.4	3.36	0.048
Resumption of job	19.73	13.2	13	<0.001
PSAS	37.93	20.6	17.64	<0.001
Patient satisfaction	112.27	113.8	114.07	0.762
Treatment cost [INR]	13963.33	14150.20	23049.86	0.003

[Table/Fig-6]: Post-operative outcomes.

[<3 days], CLA took significantly less time compared to other two procedures [OA=58.0, CLA=46.8, SPLA=79.6 minutes; p=0.027].

Significant statistical difference was noted in postoperative pain perception, measured as visual analog scale score [VASS], among three treatment arms [Table/Fig-5].

Overall mean VASS was maximum [5.05/10] at 6th hour. Mean VASS

in immediate postoperative period (first and second hour) was higher for CLA (5.07, 4.27) and SPLA (4.5, 4.79) compared to OA (2.4, 3.0) (p<0.001 and 0.001). Pattern of VASS started reversing at 4th hour when we found it to be higher for OA (5.27) than CLA (4.2) and SPLA (3.71) (p=0.001). VASS at six hours remained elevated for all treatment arms and was comparable to overall mean (5.05) (p=0.387). VASS at 12 hours and 24 hours remained high for OA (5.4 and 4.53) compared to CLA (3.67 and 3) and SPLA (3.64 and 2.86) (p<0.001 for both 12 and 24 hours). VASS at discharge and at 10th postoperative day was substantially low for all procedures (p=0.121 and 0.237). No significant difference could be established in VASS at any point of time following surgery when VASS was compared among different age-groups and between genders. Difference was significant in VASS at first postoperative hour among patients in the BMI sub-groups (6 for BMI>30, 2.8 for BMI≤18.5, p=0.007) as well as at six hours (5.4 for BMI>30, 4.65 for BMI=18.5-25, p=0.041). BMI groups were also significantly different (0 for BMI>30 and 1 for BMI≤18.5) from each other for VASS at 10th day (p=0.023).

Postoperative pain was significantly higher as the duration of surgery increased. Difference noted in mean VASS at 2nd hour was significant (6 for >120 minutes, 3.44 for 31-60 minutes; p=0.047) as was at 4th hour (4.94 for 31-60 minutes, 3.5 for 61-90 minutes, p=0.016) and 12 hours (5 for >120 minutes, 3.33 for 61-90 minutes, p=0.032). The patients who complained of pain, though mild (VASS=2), even at day 10 following surgery, had a longer duration of surgery (88 minutes against overall 65.18 minutes). No significant difference was found for comparison made between VASS at various points of time and presence of adhesions except for higher VASS at 4th postoperative hour in patients having omental adhesions (p=0.047) and at 2nd hour for patients having small bowel adhesions (p=0.006). Location of appendix did not show any bearing on VASS except for VASS at 24 hours for preileal location (six for preileal, <3.5 for others, p=0.019).

In the first hour following surgery, requirement of analgesia was significantly higher for patients who underwent either CLA or SPLA than those who underwent OA (p=0.038). Analgesia requirement again surged at 4th postoperative hour but it was now more for OA than for CLA and SPLA (p=0.072). Thereafter, in patients undergoing OA, additional analgesia continued to be needed by patients at the end of 12 (p<0.001) and 24 hours (p=0.002).

Postoperative nausea and vomiting remained less frequently occurred entity as only two patients suffered from it that too who underwent OA. Vital parameters [pulse rate, blood pressure, respiratory rate, temperature and transcutaneous oxygen saturation (SpO₂)] were recorded simultaneously with VASS and analgesia requirement. Difference found in any parameter among treatment arms or among various other groups based on age, gender, BMI or duration of surgery was not significant.

Almost 45% patients had their bowel sounds heard at two hours following surgery (OA=6, CLA=6, SPLA=8). Number of patients whose bowel sounds were present at the end of six hours rose to 75% (OA=9, CLA and SPLA each=12). Mean duration after which bowel sounds appeared in OA, CLA and SPLA was 6.27, 4.8 and 3.86 hours, respectively (p=0.336).

Wound infection remained low in the study (4, 9.1%). Out of these four patients, two patients underwent OA and one each CLA and SPLA (p=0.780). Overall mean duration of hospital stay was 3.68 days. Patients who underwent OA stayed for a significantly longer duration (OA=5.27, CLA=2.4, SPLA=3.36 days, p=0.048). Patients who underwent CLA and SPLA resumed their usual daily schedule [calculated as return to job] almost a week earlier (13.2 and 13 days, respectively) than OA (19.73 days) (p<0.001). Mean patient scar acceptance scale (PSAS) was 25.57 ranked as mostly acceptable to patients. PSAS was highest for patients who underwent OA (37.93) (CLA=20.60, SPLA=17.64, p<0.001). Scar

acceptability was no different between the two genders. Overall mean patient satisfaction score was 113.36. Patient satisfaction was almost similar in all treatment arms (OA=112.27, CLA=113.8, SPLA=114.07, p=0.762). Mean cost of treatment in OA and CLA was INR 13,963.33 and INR 14,150.20 compared to SPLA which cost INR 23049.86 (p=0.003) [Table/Fig-6].

Author	Year	Place	Sample size	Study design	Procedure
Present	2014	India	44	RCT, double blinded	OA, CLA, SPLA
Minutolo V et al., [2]	2014	Italy	230	Retrospective	OA, CLA
Sateesh et al., [3]	2014	India	50	Prospective, observational	CLA, SPLA
Frutos et al., [4]	2013	Spain	184	RCT	CLA, SPLA
Baik SM et al., [5]	2013	Korea	89	Prospective, observational	CLA, SPLA
Liang HH et al., [6]	2013	Taiwan	688	Retrospective	CLA, SPLA
Kang J et al., [7]	2012	Korea	217	RCT	CLA, SPLA
Lee JS et al., [8]	2012	Korea	63	Retrospective	CLA, SPLA
Goudhar BV et al., [9]	2011	India	204	RCT	OA, CLA
Kehagias I et al., [10]	2008	Greece	293	Non-RCT, prospective	OA, CLA
Fukami Y., [11]	2007	Japan	73	Retrospective	OA, CLA
Katkhouda N et al., [12]	2005	USA	247	RCT, double blinded	OA, CLA
De U et al., [13]	2005	India	278	Non-RCT, prospective	OA, CLA

[Table/Fig-7]: Comparison of various study designs.

Author	Duration of Surgery [Minutes]			Hospital Stay [Days]			Return to Job [Days]		
	Open	CLA	SPLA	Open	CLA	SPLA	Open	CLA	SPLA
Present	56.93	57.00	82.79	5.27	2.40	3.36	19.73	13.20	13.00
Minutolo [2]	49.3	52.2	-	3.87	2.75	-	OA>LA		-
Sateesh [3]	-	34.2	39.0	-	2.08	1.8	-	NA	NA
Baik [5]	-	71.7	71.6	-	4.5	4.3	-	NA	NA
Frutos [4]	-	32.12	38.13	-	0.78*	0.88*	-	NA	NA
Liang [6]	-	50	60	-	2.29	3	-	NA	NA
Kang [7]	-	61.7	65.88	-	3.05	2.89	-	NA	NA
Li [14]	12.35* [OA<CLA]		-	0.60*[CLA < OA]		-	4.52* [OA>CLA]		-
Kehagias [10]	47	44.3	-	3.1	2.2	-	NA	NA	-
Katkhouda [12]	60	80	-	3	2	-	No Difference		-
De [13]	25*	30*	-	5*	3*	-	14*	3*	-
Ignacio [15]	NA	NA	-	OA>CLA	0.9#	-	11 [OA=CLA]		-

[Table/Fig-8]: Duration of surgery, length of hospital stay and return to usual activity/job.

*Difference in duration; +Median, all other are mean, *Originally in hours, converted to days for comparison, Underlined values are significant.

DISCUSSION

Most of literature available regarding comparison of open and minimal invasive approach, especially SPLA is recent and whatever is available, is still sparse in finding out good study designs involving SPLA. This is further complicated by the fact that there is no standardization of SPLA, with various studies involving a unique way of doing it. A comprehensive comparison of designs of studies done in last 10 years is presented in [Table/Fig-7].

Most of the studies have involved patients with similar age groups (25-35 years) and BMI (20-26). However, gender distribution was much broad (female/male ratio 0.7-1.5).

Time taken to perform SPLA was 26 minutes more than OA and CLA in our study. However, difference in duration between CLA and OA was negligible. We finished one SPLA within 30 minutes and two cases took less than an hour. Difference in mean duration of surgery in treatment arms in patients with BMI<25 was significant but not in patients with BMI ≥25. Patients were also compared in a similar setting in reference to duration of pain [<3 days versus ≥ 3 days] and we noted that CLA took significantly less time compared to OA and SPLA in patients who presented with a shorter duration of pain. Literature is extremely diverse when it comes to time taken to finish surgery. The relatively old studies show difference between OA and CLA is more than what we see today and reflects gain of skill in doing laparoscopic procedures. The same is true for SPLA as most studies show significant difference between SPLA and other two procedures and insignificant difference between OA and CLA. A longer time taken in performing SPLA might be attributed to use of existing laparoscopic instruments, actually not designed for and therefore, surgeons faced a greater difficulty and consequently a longer operating time [Table/Fig-8].

Most studies have not compared treatment groups with respect to BMI, duration of pain, presence of adhesions, initially visualized organ, and location of appendix. So, a comparative analysis of the

Author	Postoperative pain			
	Early POP		Late POP	
	Finding	p	Finding	P
This Study	MIS>OA	<0.001	OA>MIS*	<0.001
Goudhar [9]	OA=CLA	--	OA>CLA	0.0123
Kaplan [19]	OA>CLA, <0.05			
Kehagias [10]	OA=CLA, p=0.93 [p=0.82**]			
Golub [16]	OA>CLA			
Baik [5]	SPLA>CLA	0.048	SPLA=CLA	--
Park [17]	SPLA>CLA			

[Table/Fig-9]: Post-operative Pain.

*Includes both CLA and SPLA, **Degree of pain remission.

Author	Open	CLA	SPLA
This Study	6.27 (BS)	4.80 (BS)	3.86 (BS)
Minutolo [2]	33.6 (OF)	28.4 (OF)	N/A
Baik [5]	N/A	30.6 (OF)	28.4 (OF)
Liang [6]	N/A	22 (OF)	12 (OF)
Li X et al., [14]	8 OA > CLA (OF)		N/A
Katkhouda [12]	24 (OF)	23.5 (OF)	N/A
De [13]	72 (OF)	24 (OF)	N/A

BS=Bowl Sound, OF=Oral Feeding

[Table/Fig-10]: Resumption of Oral Feeding.

present study with other studies could not be done.

Patients who underwent minimal invasive procedures in our study complained of more pain in immediate postoperative period (till 4th hour) compared to those who underwent OA. A reversal in pattern of pain from 4th hour onwards and persistence of statistically significant pain in patients in OA arm was noted at 12th and 24th hour following surgery. A detailed comparison of postoperative pain perception in various studies has been shown in [Table/Fig-9].

Pain in immediate postoperative period in laparoscopy may be explained by persistence of hypercarbia which may take some time to fully wash off from peritoneal cavity. Persistence of capnoperitoneum is a well-known entity that can cause irritation of nerve endings, operating site pain as well as referred pain elsewhere like shoulder pain [18]. We found most of the patients tolerated mild to moderate pain well up to score of 5 to 6. All the patients were offered analgesics on demand in addition to, regularly advised analgesia. Analgesia requirement was in coherence with VASS

with a higher demand reflecting an increasing VASS, followed by a subsequent decline. Requirement for additional analgesics was more for patients in CLA and SPLA than OA in early postoperative period. Requirement reversed and then persisted in favor of patients in OA arm, who demanded more analgesics after 4 hours. Kaplan, Kehagias and Baik didn't find any difference in total analgesic requirement in their respective study groups [5,10,19]. In contrast, Goudhar found requirement was more in OA arm (1.5+/-0.5, 2.5+/-0.5, p=0.3239) [9].

Incidence of PONV remained low in our study (Only 2 patients in OA, none in CLA and SPLA). Frutos has mentioned occurrence of vomiting in one patient [1.07%] in CLA [4]. Most studies have not considered vital parameters worth comparing. Resumption of oral feeds following any GI-surgery is a major milestone in recovering from surgery as well as anaesthesia. Nearly 75% patients were allowed orally after six hours, remaining were kept NPO and allowed orally next morning. Mean time difference was not significant, however, studies elsewhere showed a striking difference in duration before resumption of oral feed [Table/Fig-10].

Wound infection occurred in <10% of patients in our study. Incidence was more in OA than in CLA and SPLA but was insignificant. High incidence of wound infection in open surgeries compared to minimal

Author	Open (%)	CLA (%)	SPLA (%)
This Study	9.1	6.6	7.14
Minutolo [2]	5.49	0	-
Sateesh [3]	-	8	12
Liang [6]	-	15.7	7.1
Baik [5]	-	0	3.9
Kang [7]	-	8.5	5.3
Li [14]	8.4	3.8	-
Kehagias [10]	12.8	5.3	-
De [13]	14	4	-
Katkhouda [12]	39.1	33.3	-

[Table/Fig-11]: Wound infection.

invasive group has been supported by many studies [Table/Fig-11]. Increased wound infection in OA may be due to removal of inflamed appendix directly through the wound, whereas in LA it is extracted via a bag or trocar. In addition, port-sites in LA are smaller compared to larger wounds of OA, especially in obese patients. This result is consistent with the data shown in a recent meta-analysis, which reported a lower rate of postoperative complications, especially surgical wound infection rate, after LA [14]. Infection of the surgical wound worsens the quality of life in early postoperative period and prolongs recovery time. An advantage of LA is to reduce the wound infection rate. Mean hospital stay in our study was significantly less in CLA and SPLA than in OA [difference of >2 days] and this was similar to the findings of other reported series [Table/Fig-8]. Discrepancies in various studies may be due to social standards, insurance system and health care policies. Even in our study a relatively longer duration of hospital stay can be attributed to hilly terrain and also to the fact that most of the patients were from remote places where optimum postoperative health care facilities are still unavailable.

Return to activity following appendectomy is the subject of intense debate. A minimally invasive operation by definition should allow for a quicker recovery, shorter convalescence at home, and quicker return to work. Similar results were shown by Li and De [13,14]. In contrast, Ignacio didn't find difference in time to return to work [15] [Table/Fig-8]. Though, we found minimal difference in resumption of normal activity following CLA and SPLA, difference between OA and minimum invasive procedures was significantly more [>6 days]. These findings are supported by St Peter and a meta-analysis by Zhou [20,21]. The meta-analysis concluded days to normal activity

were significantly shorter in the SILA/SPLA than in the CLA (weighted mean difference=-0.58, 95% CI=-1.02 to -0.14, p=0.01).

CLA has been demonstrated to have advantages in certain situations such as improved cosmetic outcome. Our study showed significantly better cosmetic results in CLA than OA. Similar results were shown by Pedersen (p<0.001), De and Goudhar [9,13,22]. SPLA too showed significantly better cosmetic results than CLA. Baik and Frutos mentioned SPLA has an advantage over CLA in terms of cosmetic concerns [4,5]. Liang mentioned lower abdominal and suprapubic surgical wounds were seen more obviously in patients who underwent CLA, whereas SPLA incision scar was well hidden in the skin folds of the navel [6]. Sateesh mentioned good scar healing in SPLA, which shows better cosmetic appearance than CLA group [3].

Our analysis showed no statistical significant difference in terms of total cost of treatment between OA and CLA. Similar findings were shown by Mintulo without any statistically significant difference (p=0.812) [2]. In a meta-analysis of 8 RCT'S, Wei compared LA and OA on the basis of the cost across different countries using the hospital cost ratio and found the difference insignificant between the cost of OA and LA [23]. However, these data are in contrast with those recently published by McGrath who compared the costs between LA and OA in 2,887,823 patients undergoing surgery in the period between 1998 and 2008 [24]. Similarly the cost of treatment was higher in LA in many studies and can be attributed to the use of disposable laparoscopic instruments and the longer operative time [10,25,26]. In our study, mainly by employing reusable laparoscopic instruments, we were able to minimize the operative costs. The cost of the treatment (Rupees) was 23049 in SPLA, and 14,150 in CLA (p= 0.132). Baik showed mean cost of treatment is marginally higher in CLA as compared to SPLA [5]. In contrast Lee showed the cost was significantly lower in TULA/SPLA group; this is because the instrument was made using slim pipes and trocar [27]. Differing time under anaesthesia due to different operative time also can be one of the causes for this lack of difference in cost.

We couldn't find a significant difference in overall satisfaction of patients in the three treatment groups. Sateesh found the patient satisfactory score for CLA to be in between 6-10, with mean value of 8.04, whereas for SPLA it was in between 8-10 with mean value of 9.08 (p≤0.001) [3].

LIMITATION

We took cases of only uncomplicated appendicitis and therefore, the actual results may be different especially regarding conversion, duration of surgery, scar acceptability and hospital stay. The results may be applicable to patients in urban area where patients present early without complications but definitely not in rural area where waiting till eleventh hour still an unavoidable rule.

CONCLUSION

For lean and thin persons with BMI <25, OA is an overall better option, CLA is good for overweight and obese. All three procedures are safe. Conversion is usually not required if it's a simple appendicitis. Though immediate pain is comparable, patient do complain of prolonged postoperative pain in OA. Patients are usually satisfied in terms of scar as well as overall care, irrespective of treatment modality offered. Cost effectiveness might be an issue especially in developed countries where a large population still can't afford their basic daily requirements and not yet covered by health insurance.

REFERENCES

- [1] Amyand C. Of an inguinal rupture, with a pin in the appendix coeci, incrustrated with stone; and some observations on wounds in the guts. *Phil Trans Royal Soc.* 1736;39:329-42.
- [2] Minutolo V, Licciardello A, Stefano BD, Arena M, Arena G, Antanocci V. Outcomes and cost analysis of laparoscopic versus open appendectomy for treatment of acute appendicitis: 4 years' experience in a district hospital. *BMC Surg.* 2014; 19:14. DOI: 10.1186/1471-2482-14-14.

- [3] Sateesh S, Subhraj H, Mahesh G, Rao PS. Comparative analysis between single incision and conventional laparoscopic appendectomy for acute appendicitis. *Int J Res Med Sci.* 2014;2(4):1626-31. DOI: 10.5455/2320-6012.ijrms20141171.
- [4] Frutos MD, Abrisqueta J, Lujan J, Abellan I, Parrilla P. Randomized prospective study to compare laparoscopic appendectomy versus umbilical single-incision appendectomy. *Ann Surg.* 2013;257(3): 413-18. DOI: 10.1097/SLA.0b013e318278d225.
- [5] Baik SM, Hong KS, Kim YI. A Comparison of transumbilical single-port laparoscopic appendectomy and conventional three-port laparoscopic appendectomy: from the diagnosis to hospital cost. *J Korea Surg Soc.* 2013;85(2): 68-74. DOI: 10.4174/jkss.2013.85.2.68.
- [6] Liang HH, Hung CS, Wang W, Tam KW, Chang CC, Liu HH, et al. Single-incision versus conventional laparoscopic appendectomy in 688 patients: a retrospective comparative analysis. *Can J Surg.* 2014;57(3): E89-E97. DOI: 10.1503/cjs.023812.
- [7] Kang J, Bae BN, Gwak G, Park I, Cho H, Yang K, et al. Comparative study of a single-incision laparoscopic and a conventional laparoscopic appendectomy for the treatment of acute appendicitis. *J Korean Soc Coloproctol.* 2012;28(6):304-08. DOI: 10.3393/jksc.2012.28.6.304.
- [8] Lee JS, Choi YI, Lim SH, Hong TH. Transumbilical single port laparoscopic appendectomy using basic equipment: a comparison with the three ports method. *J Korean Surg Soc.* 2012;83(4):212-17. DOI: 10.4174/jkss.2012.83.4.212.
- [9] Goudhar BV, Telkar S, Lamani YP, Shirbur SN, Shailesh ME. Laparoscopic versus open appendectomy: a comparison of primary outcome studies from southern India. *Journal of Clinical and Diagnostic Research.* 2011;5(8):1606-09.
- [10] Kehagias I, Karamanakis SN, Panagiotopoulos S, Panagopoulos K, Kalfarentzos F. Laparoscopic versus open appendectomy: which way to go? *World J Gastroenterol.* 2008;14(31):4909-14. DOI: 10.3748/wjg.14.4909.
- [11] Fukami Y, Hasegawa H, Sakamoto E, Komatsu S, Hiromatsu T. Value of laparoscopic appendectomy in perforated appendicitis. *World J Surg.* 2007;31(1):93-97. DOI: 10.1007/s00268-006-0065-x.
- [12] Katkhouda N, Mason RJ, Towfig S, Gevorgyan A, Essani R. Laparoscopic versus open appendectomy - a prospective randomized double-blind study. *Ann of Surg.* 2005;242(3):439-450. DOI: 10.1097/01.sla.0000179648.75373.2f.
- [13] De U. Laparoscopic Versus Open Appendectomy: An Indian Perspective. *J Minim Access Surg.* 2005;1(1):15-20. DOI: 10.4103/0972-9941.15241.
- [14] Li X, Zhang, J, Sang L, Zhang W, Chu Z, Li Xin, et al. Laparoscopic versus conventional appendectomy - a meta-analysis of randomized controlled trials. *BMC Gastroenterology.* 2010;3;10:129. DOI: 10.1186/1471-230X-10-129.
- [15] Ignacio RC, Burke R, Spencer D, Bissell C, Dorsainvil C, Lucha PA. Laparoscopic versus open appendectomy: what is the real difference? Results of a prospective randomized double-blinded trial. *Surg Endosc.* 2004;18(2):334-37. DOI: 10.1007/s00464-003-8927-x.
- [16] Golub RI, Siddiqui F, Pohl D. Laparoscopic versus open appendectomy: a metaanalysis. *J Am Coll Surg.* 1998;186(5):545-53. DOI: 10.1016/S1072-7515(98)00080-5.
- [17] Park JH, Hyun KH, Park CH, Choi SY, Choi WH, Kim DJ, et al. Laparoscopic vs transumbilical single-port laparoscopic appendectomy; results of prospective randomized trial. *J Korean Surg Soc.* 2010;78:213-18. DOI: 10.4174/jkss.2010.78.4.213.
- [18] Neudecker J, Sauerland S, Neugebauer E, Bergamaschi R, Bonjer HJ, Cuschieri A, et al. The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surgical Endoscopy.* 2002;16(7):1121-43. DOI: 10.1007/s00464-001-9166-7.
- [19] Kaplan M, Salman B, Yilmaz TU, Oguz M. A quality of life comparison of laparoscopic and open approaches in acute appendicitis: a randomised prospective study. *Acta Chir Belg.* 2009;109(3):356-63. DOI: 10.1080/0015458.2009.11680439.
- [20] St Peter SD, Adibe OO, Juang D, Sharp SW, Garey CL, Laituri CA, et al. Single incision versus standard 3-port laparoscopic appendectomy: a prospective randomized trial. *Ann Surg.* 2011;254(4):586-90. DOI: 10.1097/SLA.0b013e31823003b5.
- [21] Zhou H, Jin K, Zhang J, Wang W, Sun Y, Ruan C, et al. Single incision versus conventional multiport laparoscopic appendectomy: a systematic review and meta-analysis of randomized controlled trials. *Dig Surg.* 2014;31:384-91. DOI: 10.1159/000369217.
- [22] Pedersen AG, Petersen OB, Wara P, Rønning H, Qvist N, Laurberg S. Randomized clinical trial of laparoscopic versus open appendectomy. *Br J Surg.* 2001;88(2):200-05. DOI: 10.1046/j.1365-2168.2001.01652.x.
- [23] Wei B, Qi CL, Chen TF, Zheng ZH, Huang JL, Hu BG, et al. Laparoscopic versus open appendectomy for acute appendicitis: a metaanalysis. *Surg Endosc.* 2011;25(4):1199-208. DOI: 10.1007/s00464-010-1344-z.
- [24] McGrath B, Buckius MT, Grim R, Bell T, Ahuja V. Economics of appendicitis: cost trend analysis of laparoscopic versus open appendectomy from 1998 to 2008. *J Surg Res.* 2011;171:e161-e68. DOI: 10.1016/j.jss.2011.06.067.
- [25] Kurtz RJ, Heimann TM. Comparison of open and laparoscopic treatment of acute appendicitis. *Am J Surg.* 2001;182(3):211-14. DOI: 10.1016/S0002-9610(01)00694-8.
- [26] Milewicz M, Michalik M, Ciesielski M. A prospective, randomized, unicenter study comparing laparoscopic and open treatments of acute appendicitis. *Surg Endosc.* 2003;17:1023-28. DOI: 10.1007/s00464-002-9112-3.
- [27] Lee YS, Kim JH, Moon EJ, Kim JJ, Lee KH, Oh SJ, et al. Comparative study on surgical outcomes and operative costs of transumbilical single-port laparoscopic appendectomy versus conventional laparoscopic appendectomy in adult patients. *Surg Laparosc Endosc Percutan Tech.* 2009;19(6):493-96. DOI: 10.1097/SLE.0b013e318c15493.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Surgery, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India.
2. Associate Professor, Department of Surgery, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India.
3. Associate Professor, Department of Surgery, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India.
4. Consultant Surgeon, Department of Surgery, ILS Hospital, Agartala, Tripura, India.
5. Assistant Professor, Department of Surgery, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India.
6. Senior Resident, Department of Surgery, Maharishi Valmiki Hospital, New Delhi, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Varun Kumar Singh,
Assistant Professor, Department of Surgery, Sikkim Manipal Institute of Medical Sciences,
5th Mile, Tadong, Gangtok-737102, Sikkim, India.
E-mail: varunsingh.vks@gmail.com

Date of Submission: **Dec 07, 2016**Date of Peer Review: **Mar 02, 2017**Date of Acceptance: **Jul 01, 2017**Date of Publishing: **Oct 01, 2017****FINANCIAL OR OTHER COMPETING INTERESTS:** None.