

# Variation in Facial Index and Nasal Index in Western Maharashtra Males- An Anthropometric Study

LAXMI PRASHANT NIVALE<sup>1</sup>, ACHLESHWAR GANDOTRA<sup>2</sup>, RR KARAMBALEKAR<sup>3</sup>

## ABSTRACT

**Introduction:** Facial anthropometry had proved its great importance in the field of facial reconstructive surgery as well as in the field of forensic science. Specific face types and nose types provides us important information about specific types of races.

**Aim:** To study the morphometric variation in facial index and nasal index as well as different types of faces and types of noses in the Western Maharashtra males.

**Materials and Methods:** An observational study was carried out in the five cities- Sangli, Kolhapur, Islampur, Karad and Satara of Western Maharashtra, India. Total 535 male students of different colleges between the age group of 18-20 years were selected and measurements were taken with the help of digital Vernier calliper in the sitting position after due approval from

Institutional Ethical Committees and informed consent. The methodology adopted for the measurements was taken from the guidelines given by Farkas LG in his book-“Anthropometric facial proportions in Medicine”.

**Results:** The mean facial index of Western Maharashtra males was 89.51 with standard deviation 4.26 and dominant face type was Leptoprosopic which was 46.73% and rare type of face was Hypereuriprosopic which was 0.19%. Mean nasal index of Western Maharashtra males was 72.84 with standard deviation 6.86 and dominant nose type was mesorrhine which was 63.74%.

**Conclusion:** Certain facial and nasal features are common in particular racial group. Its detection can be helpful in reconstructive facial surgeries.

**Keywords:** Anthropometry, Face types, Morphometric variation, Nose types

## INTRODUCTION

Anthropometric study of human beings had its roots in ancient days still it is used commonly all over the world as it has proven its importance in the field of cosmetic surgery, facial reconstructive surgery, and dental treatments and also in forensic science [1]. Anthropometric studies are scientific methods and techniques for displaying different measurement and observation on the human being as well as skeleton. Anthropometric studies are very important area for craniofacial surgery and syndromology [2]. Study of nose is also useful in the determination of race and sex of individual or group whose identity is unknown [3]. Facial index in general and nasal index in particular are useful in forensic investigations [4]. Facial index and nasal index provides us an important information about race and ethnicity. By knowing nasal index and type of nose it can be concluded about climatic condition. According to theory of natural selection, people who lives in cold and dry weather possesses long and narrow type of nose and who lives in warm and moist weather possesses broad nose [5]. With this background, the present study was conducted with an aim to study different facial and nasal parameters in adult male population of Western Maharashtra, India.

## MATERIALS AND METHODS

This was an observational type of study carried out from March 2018 to January 2020 over 535 male students of following colleges as shown in [Table/Fig-1]. After getting due approval from both the Institutes: Prakash Institute of Medical Sciences and Research, Urun-Islampur (Ref No.-PSM/PIMSR/24013/2018) and Sumandeep Vidyapeeth University, Vadodara, Gujarat, India (SVIEC/IN/MEDI/PHD/18004) the study was conducted. NOC letter was obtained from Dean/Principal of respective college. The procedure and purpose of the study was explained to the participants and then written consent was obtained from them.

| Sr. No. | Region   | College name  | No. of students |
|---------|----------|---|-----------------|
| 1       | Islampur | Prakash Institute of Medical Sciences and Research, Urun- Islampur                            | 93              |
|         |          | Loknete Rajaram Bapu Patil Ayurvedic Medical College  | 30              |
| 2       | Kolhapur | Dr JJ Magdum College of Engineering, Jaysingpur   | 52              |
|         |          | Dr JJ Magdum Homoeopathic Medical College, Jaysingpur   | 27              |
|         |          | Dr JJ Magdum Ayurvedic Medical College, Jaysingpur  | 22              |
| 3       | Sangli   | Hon. Shri Annasaheb Dange Ayurvedic Medical College, Post Graduate and Research Centre, Ashta | 38              |
|         |          | Hon. Shri Annasaheb Dange College of Pharmacy (B.Pharm)                                       | 46              |
|         |          | Hon. Shri. Annasaheb Dange College of Pharmacy (D.Pharm)                                      | 27              |
| 4       | Satara   | Gourishankar Institute of Pharmaceutical Education and Research, Limb                         | 45              |
|         |          | Mahalaxmi Institute of Pharmacy (D. Pharm), Raigaon   | 28              |
|         |          | Late. Narayandas Bhawandas Chhabada Institute of Pharmacy, Raigaon                            | 25              |
| 5       | Karad    | Krishna Institute of Medical Sciences “Deemed To Be University” Karad                         | 84              |
|         |          | Krishna Institute of Nursing Science, Karad   | 18              |


[Table/Fig-1]: Number of students recruited from different cities and colleges.

All the participants were residing in the respective city from their forefathers, after taking personal and family history in brief with an age range of 18-20 years were included. This age group was selected because development of face has been completed at 18 years. Students who were not resident of Western Maharashtra

were excluded from the study. Individuals with normal craniofacial configuration were selected and those who were having background of genetically transmitted disorders like cleft lip, cleft palate were also excluded from the study. Calibrated Standard Vernier calliper (30 centimetre length with accuracy of 0.01 mm) was used to take facial measurements. All the facial soft tissue landmarks were first marked with the help of skin marking pencil and then the measurements were taken in sitting position by the author by standing in front of the participants. Keeping the left hand on the head of the subject and holding the tip of upper arm of the calliper with thumb and index finger on the marked point of face and sliding the lower arm of calliper in upward direction up to desired marking so that the tip just touches the desired point. Participants were asked to remain quiet with erect neck with gently closed lips and avoid talking or laughing while taking measurements. All the facial measurements were taken in millimetres.

The following facial measurements were taken-

2a) Facial Height/length (n-mn/gn)- distance between nasion (n) to menton (mn/gn); 2b) Facial width/breadth- distance between right and left zygion (zy-zy); 2c) Height/Length of nose (n-sn)- distance between nasion (n) to subnasale (sn); 2d) Width of nose (al-al)- distance between right and left alar of nose [Table/Fig-2a-d].



**[Table/Fig-2]:** a) Showing measurement of face height (n-mn); b) Showing measurement of face width (zy-zy); c) Shows height of nose (n-sn); d) Shows width of nose (al-al).

**Facial index:** Face length (n-mn/gn)/Face width (zy-zy)×100.

It is called as prosopic facial index and different face types were categorised according to Banister classification [6] in the text book of medical anthropometry by Farkas LG in Anthropometric facial proportion in Medicine [7] which are as follows:

- 1) Hypereuriprosopic (very broad face)- when the facial index is lesser than or equal to 79.9 [Table/Fig-3].
- 2) Euriprosopic (broad face)- when the facial index is between 80 to 84.9 [Table/Fig-4].
- 3) Mesoprosopic (round face)- when the facial index is between 85 to 89.9 [Table/Fig-5].
- 4) Leptoprosopic (long face)- when the facial index is between 90 to 95 [Table/Fig-6].
- 5) Hyperleptoprosopic (very long face)- when the facial index is above 95 [Table/Fig-7].



**[Table/Fig-3-7]:** Different types of faces. (Image from left to right)

**Nasal index:** Width of nose (al-al)/Length of nose (n-sn)×100.

Paul Topinard (1830-1911) defined the nasal index [8]. On the basis of this index, the following three types of noses were categorised:

- 1) Leptorrhine (long nose)- when the nasal index <70.
- 2) Mesorrhine (medium nose)- when the nasal index is between 70 to 84.9.
- 3) Platyrrhine (flat nose)- when the nasal index is >85.

## STATISTICAL ANALYSIS

All the collected data were subjected to appropriate statistical analysis (mean, standard deviation). It was analysed by using descriptive statistics. Software used was R statistical software with 3.5 version.

## RESULTS

Mean face height (142.54 mm), face width (125.85 mm), nose height (52.76 mm) and nose width (38.27 mm) are shown in [Table/Fig-8] with standard deviation. The study results showed that mean facial index in Western Maharashtra males was 89.51 with standard deviation 4.26 and mean nasal index was 72.84 with standard deviation 6.86 [Table/Fig-9].

| Variable    | Mean (mm) | Standard deviation |
|-------------|-----------|--------------------|
| Face length | 142.54    | 6.68               |
| Face width  | 125.85    | 5.52               |
| Nose length | 52.76     | 4.04               |
| Nose width  | 38.27     | 2.72               |

**[Table/Fig-8]:** Mean- facial length, facial width, nose length, nose width with standard deviation.

| Variable     | Mean  | Standard deviation | Minimum | Maximum |
|--------------|-------|--------------------|---------|---------|
| Facial index | 89.51 | 4.26               | 89.47   | 91.39   |
| Nasal index  | 72.84 | 6.86               | 55.74   | 89.74   |

**[Table/Fig-9]:** Mean, standard deviation, maximum and minimum values for facial and nasal index.

The dominant type of male face was leptoprosopic (46.73%). Similarly, the dominant nose type was mesorrhine (63.74%) followed by leptorrhine (34.02%) [Table/Fig-10,11].

| Type of face       | No. of subjects | Percent |
|--------------------|-----------------|---------|
| Euriprosopic       | 103             | 19.25%  |
| Hypereuriprosopic  | 1               | 0.19%   |
| Hyperleptoprosopic | 14              | 2.62%   |
| Leptoprosopic      | 250             | 46.73%  |
| Mesoprosopic       | 167             | 31.21%  |
| Total              | 535             | 100.00% |

**[Table/Fig-10]:** Distribution of face types in Western Maharashtra males.

| Type of nose | No. of subjects | Percent |
|--------------|-----------------|---------|
| Leptorrhine  | 182             | 34.02%  |
| Mesorrhine   | 341             | 63.74%  |
| Platyrrhine  | 12              | 2.24%   |
| Total        | 535             | 100%    |

**[Table/Fig-11]:** Distribution of nose types in Western Maharashtra males.

## DISCUSSION

The present study results showed the predominant face type of Western Maharashtra males was leptoprosopic. It correlates with study done in South Indian and North Indian population by Ashwini C et al., [9]. It is also in conformity with Uttarakhand males by Ansari S et al., [10]. It contradicts with males in Haryana by Anand S et al., and with Gujarati males by Kanan U et al., [11,12].

From [Table/Fig-12], it was clear that Serbian male (Jeremic D et al.) showed similar face type with Western Maharashtra males while Nepali males (Pandey N et al.), Malay males (Yesmin T et al.) and Turkish males (Ozsahm E et al.) were having different face type than males in present study [9-16].

| Name of Author                       | Population          | Hypereuriprosopic | Euriprosopic | Mesoprosopic | Leptoprosopic | Hyperleptoprosopic |
|--------------------------------------|---------------------|-------------------|--------------|--------------|---------------|--------------------|
| Ashwini C and Karinaganannavar A [9] | South India         | 9.09              | 18.18        | 10.9         | 40            | 21.81              |
|                                      | North India         | 0                 | 14.28        | 28.57        | 46.42         | 10.71              |
| Ansari S et al., [10]                | Uttarakhand         | 4                 | 11           | 28           | 37            | 20                 |
| Anand S et al., [11]                 | Haryana             | 11                | 43           | 53           | 27            | 16                 |
| Kanan U et al., [12]                 | Gujarat             | 35.2              | 43           | 18.2         | 3.6           | 0                  |
| Pandey N et al., [13]                | Nepal               | 13                | 23           | 48.66        | 13.33         | 2                  |
| Jeremic D et al., [14]               | Serbia              | 0                 | 0            | 17.78        | 76.67         | 5.56               |
| Yesmin T et al., [15]                | Malay               | 12                | 18           | 45           | 20            | 5                  |
| Ozsahm E et al., [16]                | Turkish             | 18.1              | 35.36        | 33.21        | 8.7           | 4.7                |
| Present study                        | Western Maharashtra | 0.19              | 19.25        | 31.21        | 46.73         | 2.65               |

**[Table/Fig-12]:** Shows comparison of male face types with other studies in India in percentage [9-16].

Present study showed the mean nasal index in Western Maharashtra males was 72.84 and predominant nose type was mesorrhine. It correlates with the males of Sindhis in Rajasthan (Choudhary A and Choudhary DS), of Jammu and Kashmir (Jabeen N et al.), of Karnataka and Kerala (Shivanand ND et al.), of Uttar Pradesh (Ray SK et al.), and Santhals of West Bengal (Shah MRI et al.) in India whereas contradicts Jats in Rajasthan (Choudhary and Choudhary DS) [2] and Bengalis in West Bengal (Shah MRI et al.) in India [Table/Fig-13] [2,4,17-21].

| Author/Year                             | Population          | Nasal index | Predominant type of nose |
|---|---------------------|-------------|--------------------------|
| Shivanand ND et al., (2016) [17]        | Karnataka           | 73.37       | Mesorrhine               |
|   | Kerala              | 82.99       | Mesorrhine               |
| Jabeen N et al., (2019) [4]             | Jammu and Kashmir   | 72.15       | Mesorrhine               |
| Ray SK et al., (2016) [18]              | Uttar Pradesh       | 75.86       | Mesorrhine               |
| Choudhary A and Choudhary DS (2012) [2] | Jats                | 68.09       | Leptorrhine              |
|   | Sindhis             | 70.72       | Mesorrhine               |
| Shah MRI et al., (2015) [19]            | Santhals            | 80.00       | Mesorrhine               |
|   | Bengalis            | 65.90       | Leptorrhine              |
| Mohammad I et al., (2018) [20]          | Nigeria             | 74.08       | Mesorrhine               |
| Asthuta AR and Pradipta IPY (2019) [21] | Indonesia           | 74.08       | Platyrrhine              |
| Present study                           | Western Maharashtra | 72.84       | Mesorrhine               |

**[Table/Fig-13]:** Shows comparison of nose types with other studies [2,4,17-21].

The nose of Western Maharashtra males correlates with Nigerian males (Mohammad I et al.) but different from Indonesian male nose type Asthuta AR and Pradipta IPY [20,21].

From the above results it is clear that face types and nose type helps to determine the characteristic racial features. Present study results will help surgeons in facial reconstructive surgery and so, also in rhinoplasty of males.

## Limitation(s)

As the study is focussed on only five cities of Western Maharashtra, the results of it cannot be generalised to whole of Western Maharashtra. More extensive work is required for generalisation.

## CONCLUSION(S)

Certain human facial features are commonly found among certain racial groups or in any specific geographical place. By finding out such a specific face type will help in reconstructive and facial plastic surgeries. Furthermore, extensive study is required in this field to establish all the parameters of face. From this study it is concluded that majority of Western Maharashtra males were having leptoprosopic type of face 46.73% and mesorrhine type of nose (63.74%).

## REFERENCES

- 1] Vegter F, Hage JJ. Clinical anthropometry and canons of the face in historical perspective. *Plastic and Reconstructive Surgery*. 2000;106(5):1090-96.
- 2] Choudhary A, Choudhary DS. Comparative anthropometric study of nasal parameters between two ethnic groups of Rajasthan state. *International Journal of Medicine and Public Health*. 2012;2(2):46-48.
- 3] Oladipo G, Udoaka A, Afolabi E, Manuel BI. Nasal parameters of Itsekiris and Urhobos of Nigeria. *Int J Biol Anthropol*. 2008;3:01-05.
- 4] Jabeen N, Magotra R, Choudhary S, Sharma AK. Study of nasal index in different zones of Jammu and Kashmir. *JK SCIENCE*. 2019;22(2).
- 5] Hall RL, Hall DA. Geographic variation of native people along the Pacific Coast. *Hum Biol*. 1995;67(3):407-26.
- 6] Banister M, Williams P, Dyson M, Dussak JE. *Gray's Anatomy*, 38<sup>th</sup> ed, London: Churchill Livingstone; 1995: 607-12.
- 7] Farkas LG, Munro IR. *Anthropometric facial proportions in medicine*. Charles Thomas Springfield: 1987.
- 8] Topinard P. *Elements d'Anthropologie Generale*, Paris. Delahaye & Lecrosnier; 1885: 244-307.
- 9] Ashwini C, Karinaganannavar A. A study of facial index among North and South Indian students of JSS Medical College. *International Journal of Life Sciences Biotechnology and Pharma Research*. 2014;3(2).
- 10] Ansari S, Singala M, Ravi KS. Facial anthropometry in adult Jaunsari tribe population of Dehradun district of Uttarakhand. *Journal of Clinical and Diagnostic Research*. 2019;13(4):AC01-AC03.
- 11] Anand S, Tripathi S, Chopra A, Khaneja K. Vertical and horizontal proportions of the face and their co-relation to phi among Indians in Moradabad population: A survey, *The Journal of Indian Prosthodontic Society*. 2015;15(2):125.
- 12] Kanan U, Gandotra A, Desai A, Andani R. Variation in facial index of Gujarati males- A photometric study. *International Journal of Medical and Health Sciences*. 2012;1(4):27-31.
- 13] Pandey N, Gogoi P, Budathoki D, Gopal KC. Anthropometric study of facial index of medical students. *Journal of Kathmandu Medical College*. 2015;4(14).
- 14] Jeremic D, Kocic S, Vulovic M, Sazdanovic M, Sazdanovic P, Jovanovic B, et al. Anthropometric study of the facial index in the population of central Serbia. *Arch Biol Sci Belgrade*. 2013;65(3):1163-68.
- 15] Yesmin T, Thwin SS, Urmi SA, Wai MM, Zaini PF, Azwan K. A study of facial index among Malay Population. *Journal of Anthropology*. 2014;2014:726974.
- 16] Ozsahm E, Kizilkanat E, Boyan N, Soames R, Oguo O. Evaluation of face shape in Turkish Individuals. *Int J Morphol*. 2016;34(3):904-08.
- 17] Shivanand ND, Bidarkotimath S, Kumar S. Comparison of nasal index in Karnataka and Kerala population. *Journal of Evidence Base of Medicine and Health Care*. 2016;3(20):823-24.
- 18] Ray SK, Saha K, Kumar A, Banjare S. Anthropometric study of nasal index among the population of Western Uttar Pradesh Region. *International Journal Scientific Study*. 2016;4(2):65-70.
- 19] Shah MRI, Anwar S, Mandal DK, Yesmin S, Ahmad S. Anthropometry of the nose: A comparative study between adult male Santhals and Bengalis in Bangladesh. *Mediscope*. 2015;2(2):28-32.

- [20] Mohammad I, Mokhtari T, Ijaz S, Omotosha AD, Ngaski AA, Milaniford M, et al. Anthropometric study of nasal index in Hausa ethnic population of northwest Nigeria. *Contemp Med Sci.* 2018;4(1):26-29.
- [21] Asthuta AR, Pradiptha IPY. Anthropometric study of nasal index of the Bali aga population. *ORLI, Tahun.* 2019;49(1):35-39.

**PARTICULARS OF CONTRIBUTORS:**

1. Ph.D Scholar, SBKS Medical College, Sumandeep Vidyapeeth Vadodara, Gujarat, India; Assistant Professor, Department of Anatomy, Prakash Institute of Medical Science and Research, Urun-Islampur, Maharashtra, India.
2. Professor and Head, Department of Anatomy, SBKS Medical College, Sumandeep Vidyapeeth University, Vadodara, Gujarat, India.
3. Professor and Head, Department of Anatomy, Prakash Institute of Medical Science and Research, Urun-Islampur, Maharashtra, India.

**NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Laxmi Prashant Nivale,  
Ph.D Scholar, SBKS Medical College, Sumandeep Vidyapeeth,  
Vadodara-391760, Gujarat, India.  
E-mail: pnivale@rediffmail.com

**PLAGIARISM CHECKING METHODS:** [Jain H et al.]

- Plagiarism X-checker: Sep 16, 2020
- Manual Googling: Oct 27, 2020
- iThenticate Software: Oct 27, 2020 (10%)

**ETYMOLOGY:** Author Origin**AUTHOR DECLARATION:**

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Sep 14, 2020**Date of Peer Review: **Oct 01, 2020**Date of Acceptance: **Oct 27, 2020**Date of Publishing: **Nov 01, 2020**