

Role of Glycaemic Control in Predicting Severity of Illness in COVID 19 Patients with Diabetes Mellitus

JAYANTHI HARI KISHAN¹, VOLLALA SHRAVAN KUMAR²

ABSTRACT

Introduction: Coronavirus Disease 2019 (COVID 19) pneumonia is a rapidly spreading disease and which causes morbidity and mortality of many patients. Diabetes mellitus is co-morbidity which is considered as the risk factor for COVID 19. Well-controlled diabetes is associated with better outcomes than poorly controlled diabetes. Measurement of glycated haemoglobin (HbA1c) is the standard method for assessing long term glycaemic control. Regardless of the level of hyperglycaemia, improvement in glycaemic control will lower the risk of diabetic complications.

Aim: This study was conducted to identify the role of glycaemic control (HbA1c) in predicting the severity of illness in patients with COVID 19 pneumonia.

Materials and Methods: This was a retrospective observational study of (51 diabetic and 51 were non diabetic) patients at Kamineni Academy of Medical Sciences, Hyderabad, India. The patients diagnosed with COVID 19 pneumonia, which

includes both diabetics and non diabetics from June 2020 to September 2020. Patients age, sex, baseline HbA1c levels, and oxygen requirement during the hospital stay were analysed using Statistical Package for the Social Sciences (SPSS) software version 22.0. The Chi-Square test was used to analyse qualitative data and p-value significant at level <0.05.

Results: In the study among diabetics (n=51), 20 (39.2%) were on room air, 24 (47.1%) required intermittent oxygen support, 3 (5.9%) high flow oxygen, and 4 (7.8%) non invasive ventilator support. Among non diabetics (n=51), 28 (54.9%) were on room air, 18 (35.3%) on intermittent oxygen, 2 (3.9%) high flow oxygen, and 3 (5.9%) Non Invasive Ventilator (NIV) support. It was observed that patients with HbA1c measurements with poor glycaemic control required more oxygen support during treatment in diabetics (p-value:0.469)

Conclusion: In the present study, patients with poor glycaemic control required insignificantly, more oxygen support than patients with good glycaemic control.

Keywords: Body mass index, Coronavirus 2019, Food preferences, Junk food, Undernutrition

INTRODUCTION

Coronavirus disease (COVID 19) is the worst pandemic in the past 100 years and has affected millions of people in more than 200 countries. The novel severe acute respiratory syndrome coronavirus (SARS-COV-2) was identified as the causative agent. A greater risk of COVID 19 has been reported in patients with diabetes and hypertension. Comparatively, the SARS epidemic in 2003, also caused by a beta coronavirus, was associated with a 3-fold risk of poor outcomes in the presence of diabetes, the highest among all comorbidities [1].

Diabetes mellitus, as a pre-existing health condition is a chronic low grade inflammation, which causes a dysregulated immune response, microvascular injury with a procoagulant state. One study of 191 patients reported a mortality risk of 2.85 fold for patients with diabetes [2]. Indeed, in patients with diabetes the insulin resistance, poor glycaemic control leads to alveolar capillary microangiopathy and interstitial fibrosis via over inflammation. A systematic review by Huang I et al. revealed that diabetes was associated with mortality, severity, acute respiratory distress syndrome, and disease progression in patients with COVID 19 [3]. In COVID 19, dichotomous outcomes of disease severity such as ICU admission and mortality were significantly elevated (~2- and ~3-fold, respectively) in diabetes. This is of particular interest because Huang I et al., noted an increased risk of a composite poor outcome in patients with diabetes, which included severe COVID 19 as one of the outcomes. One study of 1382 COVID 19 patients with diabetes found a 2.79-fold risk of admission to the ICU [4].

The present study aimed to find the association between diabetes as a co-morbidity and their pre-existing glycaemic control with clinical course of disease in COVID 19 patients and their oxygen requirement during the treatment.

MATERIALS AND METHODS

This single centre retrospective observational study was conducted from June 2020 to September 2020 on (51 diabetic and 51 non diabetic) patients, at Kamineni Academy of Medical Sciences, Hyderabad, India. The record of the patients diagnosed with COVID 19 which includes diabetic and nondiabetic patients and data, including age, sex, history of diabetes, HbA1c, level of illness severity (oxygen requirement) were noted. diagnosed with COVID 19 which includes diabetic and non diabetic patients and data, including age, sex, history of diabetes, HbA1c, level of illness severity (oxygen requirement). All the subjects in the complete article were called Diabetics and non-diabetics according to their history of Diabetes. also the HbA1c of all subjects mentioned and tabulated in results gave us a history of glycaemic control for past three months for all of them.

Inclusion criteria: Data of Real-time RT-PCR-Reverse transcription polymerase chain reaction (RT-PCR) positive patients, with type-II diabetes mellitus and non diabetics of age group >30 years was included in the study.

Exclusion criteria: COVID 19 RT-PCR negative patients data was excluded from the study.

STATISTICAL ANALYSIS

Data were entered into a Microsoft Excel datasheet and were analysed using SPSS version 22.0 (IBM SPSS Statistics, Somers NY, USA) software. Categorical data was represented in the form of frequencies and proportions in MS Excel and MS word. Chi-square test was used as a test of significance for qualitative data and p-value <0.05 was considered as statistically significant.

RESULTS

Among diabetics, the majority of subjects, 16 (31.4%) were in the age group 50 to 59 years and among non diabetics majority, 14 (27.5%) were in the age group 40 to 49 years. There was no significant difference in age distribution between diabetics and non diabetics [Table/Fig-1].

Among diabetics, 39 (76.5%) were males and 12 (23.5%) were females and among non diabetics, 37 (72.5%) were males and 14 (27.5%) were females. There was no significant difference in sex distribution between diabetics and non diabetics patients. [Table/Fig-1].

Demographic Variables		Diabetes						p-value
		Yes		No		Total		
		Count	%	Count	%	Count	%	
Age (years)	30-39	4	7.8%	12	23.5%	16	15.7%	0.053
	40-49	9	17.7%	14	27.5%	23	22.7%	
	50-59	16	31.4%	13	25.5%	29	28.4%	
	60-69	7	13.7%	6	11.7%	13	12.7%	
	>70	15	29.4%	6	11.8%	21	20.6%	
Sex	Male	39	76.5%	37	72.5%	76	74.5%	0.650
	Female	12	23.5%	14	27.5%	26	25.5%	

[Table/Fig-1]: Profile of subjects in the study.

Among diabetics, 4 (7.8%) had normal HbA1c, 9 (17.6%) had Pre-diabetic range and 38 (74.5%) had Diabetic Range >6.5% [Table/Fig-2].

HbA1c measurements	Diabetic patients	
	Count (n)	Percentage (%)
Normal < 5.7%	4	7.8%
Prediabetic range 5.7%- 6.5%	9	99.9%
Diabetic range >6.5%	38	74.5%

[Table/Fig-2]: HbA1c distribution among diabetics.


In the study among diabetics, 20 (39.2%) required room air, 24 (47.1%) intermittent oxygen, 3 (5.9%) high flow oxygen, and 4 (7.8%) NIV support. Among non diabetics, 28 (54.9%) required room air, 18 (35.3%) intermittent oxygen, 2 (3.9%) high flow oxygen, and 3 (5.9%) NIV support. There was no significant difference in oxygen support between diabetics and non diabetics. (p-value: 0.469) [Table/Fig-3,4].

Oxygen Support Data of subjects	Diabetes					
	Yes		No		Total	
	Count (n)	Percentage (%)	Count (n)	Percentage (%)	Count (n)	Percentage (%)
Room Air	20	39.2%	28	54.9%	48	47.1%
Intermittent Oxygen	24	47.1%	18	35.3%	42	41.2%
High Flow Oxygen	3	5.9%	2	3.9%	5	4.9%
NIV Support	4	7.8%	3	5.9%	7	6.9%
Total	51	100.0%	51	100.0%	102	100.0%

[Table/Fig-3]: Oxygen support comparison between diabetics and non diabetic patients ($\chi^2=2.53$, differential factor=3, p=0.469).

DISCUSSION

The prevalence of diabetes mellitus is anticipated to increase substantially during the next decades worldwide and considered to be the main cause of human deaths [5]. People with diabetes are more susceptible to certain infectious diseases, such as Staphylococcus aureus and Mycobacterium tuberculosis, possibly because of their dysregulated immune system [6,7]. Hyperglycaemia can impair host defences and poor glycaemic control has been associated with



[Table/Fig-4]: Bar diagram showing Oxygen support comparison between diabetics and non diabetics patients.

infections. Given that glycaemic control is a modifiable factor and can be achieved and sustained by health care interventions.

It is known clearly that the interplay between COVID 19 and diabetes mellitus entails complex pathophysiology. The surface proteins of the SARS-CoV-2 virus could attack heme on the 1- β chain of haemoglobin in red blood cells of individuals with diabetes, dissociating iron to form porphyrin, thereby causing less and less haemoglobin that can carry oxygen, and carbon dioxide ultimately leading to symptoms of respiratory distress [8-10]. It is suggested that deoxy haemoglobin is more vulnerable than oxidised haemoglobin to the surface proteins of the SARS-CoV-2 virus, and glycosylated haemoglobin (deoxygenated form) levels are higher in diabetic patients [8]. These findings suggest that patients with hyperglycaemia or diabetes may be more vulnerable to SARS-CoV-2 attack due to enhanced glycation of haemoglobin, thus increasing the risk for COVID 19 associated mortality rate.

The pathogenic mechanism of how the SARS-CoV-2 virus binds to the 1- β chain of porphyrins of the erythrocytes, leading to release of iron and disturbance of heme-metabolism, remains elusive and needs further investigation.

Diabetes is well known to induce pulmonary dysfunction, such as reducing lung volumes and compliance and increasing airway resistance, which is related to insulin resistance and nonenzymatic glycosylation of lung proteins [11]. This may be another risk factor for more-severe illness in COVID 19, the SARS-CoV-2 infection could cause hyper-inflammation in cells, resulting in excessive activation of macrophages, which can suppress the recruitment of T cells [12]. Emerging evidence shows that during the acute phase of infection, SARS-CoV-2 invades CD4+ T and CD8+ T lymphocytes, leading to cell apoptosis and lymphocytopenia [13]. It has been shown that elevated glucose levels promote SARS-CoV-2 replication in monocytes, resulting in inhibition of T cell response [14]. Diabetes is well known to be associated with dysfunctional innate and adaptive immunity [15] and it is thus speculated that impaired immune function in diabetes patients may accentuate SARS-CoV-2 infection and its harmful function.

Recent researches revealed that cytokine storm emerges in COVID 19 patients with hyperglycaemia or Type 2 Diabetes Mellitus (T2DM), leading to dysregulation of immune response and Acute Respiratory Distress Syndrome (ARDS) [16].

Increased levels of chronic inflammatory factors, such as IL-1 β , IL-6, TNF- α , and activation of the immune response are frequently

found in diabetic patients with poor-controlled blood glucose levels [17]. Thus, high blood glucose may promote cytokine storm emerge and immune over-activation, ultimately causing ARDS and multiple organ failure in COVID 19 patients.

It may therefore be more appropriate to say that complications and co-morbidities linked to diabetes are associated with a higher mortality rate than the presence of diabetes per se [18].

Limitation(s)

Smaller sample size and the statistical significance of the above data has to be interpreted with caution.

CONCLUSION(S)

The outbreak of the COVID 19 pandemic has caused a global health crisis of our time. A growing body of clinical evidence shows that patients with pre-existing diabetes are highly susceptible to SARS-CoV-2 infection and its associated mortality. The exact mechanisms linking diabetes and COVID 19 remain to be further elucidated, but our observations suggest that poor glycaemic control may play critical role in the severity of COVID 19 in patients with pre-existing diabetes and showed higher oxygen requirement as compared to diabetics with good glycaemic control. The requirement for further studies involving a larger sample size needs to be considered to draw further conclusions.

REFERENCES

- [1] Booth CM, Matukas LM, Tomlinson GA, Rachlis AR, Rose DB, Dwosh HA, et al. Clinical features and short-term outcomes of 144 patients with SARS in the greater Toronto area. *JAMA*. 2003;289(21):2801-09.
- [2] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID 19 in Wuhan, China: A retrospective cohort study [published correction appears in *Lancet*. 2020;395(10229):1038. [published correction appears in *Lancet*. 2020;395(10229):1038]. *Lancet*. 2020;395(10229):1054-62.
- [3] Huang I, Lim MA, Pranata R. Diabetes mellitus is associated with increased mortality and severity of disease in COVID 19 pneumonia - A systematic review, meta-analysis, and meta-regression. *Diabetes Metab Syndr*. 2020;14:395-403. doi: 10.1016/j.dsx.2020.04.018
- [4] Roncon L, Zuin M, Rigatelli G, Zuliani G. Diabetic patients with COVID 19 infection are at higher risk of ICU admission and poor short-term outcome. *J Clin Virol*. 2020;127:104354. doi: 10.1016/j.jcv.2020.104354
- [5] Knapp S. Diabetes and infection: is there a link?—a mini-review. *Gerontology*. 2013;59:99-104. doi: 10.1159/000345107
- [6] Muller LM, Gorter KJ, Hak E, Goudzwaard WL, Schellevis FG, Hoepelman AI, et al. Increased risk of common infections in patients with type 1 and type 2 diabetes mellitus. *Clin Infect Dis*. 2005;41:281-88. doi: 10.1086/431587
- [7] Hodgson K, Morris J, Bridson T, Govan B, Rush C, Ketheesan N. Immunological mechanisms contributing to the double burden of diabetes and intracellular bacterial infections. *Immunology*. 2015;144:171-85. doi: 10.1111/imm.12394
- [8] Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and mortality in adult COVID 19 inpatients in Wuhan. *The Journal of Allergy and Clinical Immunology*. 2020;146(1):110-18.
- [9] Liu WZ, Li HL. COVID 19-Attacks the 1-β chain of hemoglobin and captures the porphyrin to inhibit human heme metabolism. *ChemRxiv*. 2020. Preprint.
- [10] Means C. Letter to the Editor: Mechanisms of increased morbidity and mortality of SARS-CoV-2 infection in individuals with diabetes: what this means for an effective management strategy. *Metabolism*. 2020;108:154254. doi: 10.1016/j.metabol.2020.154254
- [11] Lecube A, Simo R, Pallayova M, Punjabi NM, Lopez Cano C, Turino C, et al. Pulmonary function and sleep breathing: two new targets for type 2 diabetes care. *Endocr Rev*. 2017;38(6):550-73.
- [12] Iacobellis G. COVID 19 and diabetes: Can DPP4 inhibition play a role? *Diabetes Research and Clinical Practice*. 2020;162:108125. doi: 10.1016/j.diabres.2020.108125
- [13] Angelidi AM, Belanger MJ, Mantzoros CS. COVID 19 and diabetes mellitus: What we know, how our patients should be treated now, and what should happen next. *Metabolism: Clinical and Experimental*. 2020;107:154245.
- [14] Codo AC, Davanzo GG, Monteiro LB, de Souza GF, Muraro SP, Virgilio-da-Silva JV, et al. Elevated glucose levels favor SARS-CoV-2 infection and monocyte response through a HIF-1α/glycolysis-dependent axis. *Cell Metab*. 2020;32(3):437-46.e5.
- [15] Donath MY, Dinarello CA, Mandrup-Poulsen T. Targeting innate immune mediators in type 1 and type 2 diabetes. *Nat Rev Immunol*. 2019;19(12):734-46.
- [16] Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID 19. *The Journal of Infection*. 2020;80(6):607-13.
- [17] Apicella M, Campopiano MC, Mantuano M, Mazoni L, Copelli A, Del Prato S, et al. COVID 19 in people with diabetes: understanding the reasons for worse outcomes. *The Lancet. Diabetes & Endocrinology*. 2020;8(9):P782-92.
- [18] Guo W, Li M, Dong Y, Zhou H, Zhang Z, Tian C, et al. Diabetes is a risk factor for the progression and prognosis of COVID 19. *Diabetes Metab Res Rev*. 2020.36(7) e3319. <https://doi.org/10.1002/dmrr.3319>.
- [19] Liu Z, Li J, Huang J, Guo L, Gao R, Luo K, et al. Association between diabetes and COVID 19: A retrospective observational study with a large sample of 1,880 cases in Leishenshan Hospital, Wuhan. *Front Endocrinol*. 2020;11:478. doi: 10.3389/fendo.2020.00478.

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of General Medicine, Kamineni Academy of Medical Sciences, Hyderabad, Telangana, India.
2. Postgraduate Student, Department of General Medicine, Kamineni Academy of Medical Sciences, Hyderabad, Telangana, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Vollala Shravan Kumar,
H-No 9-6-129/2 Ramna Gar, Karimnagar, Telangana-505001, India.
E-mail: sravancaims@gmail.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: No
- Was Ethics Committee Approval obtained for this study? No
- Was informed consent obtained from the subjects involved in the study? No
- For any images presented appropriate consent has been obtained from the subjects. NA

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Dec 14, 2020
- Manual Googling: Jan 11, 2021
- iThenticate Software: Mar 30, 2021 (25%)

ETYMOLOGY: Author Origin

Date of Submission: Dec 05, 2020

Date of Peer Review: Jan 09, 2021

Date of Acceptance: Feb 12, 2021

Date of Publishing: May 01, 2021