

Outcome of COVID-19 in Patients Requiring Haemodialysis- A Retrospective Observational Study

GIRISH P VAKRANI¹, TANUJA NAMBAKAM²

ABSTRACT

Introduction: Acute Kidney Injury (AKI) and Chronic Kidney Disease (CKD) are risk factors for COVID-19 infection. Patients with kidney disease also have other comorbidities like hypertension, diabetes mellitus, cardiovascular disease that are risk factors for poor outcome in COVID-19 infections. Severe COVID-19 has multiorgan involvement including AKI. Compared to normal population, renal failure patients with COVID-19 have extensive lung involvement, need more ventilator support, and have higher mortality.

Aim: To assess outcome (mortality, recovery) and correlation between factors (age, gender, co-morbidities), biochemical parameters with mortality in COVID-19 patients requiring haemodialysis.

Materials and Methods: It was a retrospective observational study wherein, data was collected, entered and analysed with

Statistical Package for the Social Sciences (SPSS) version 19.0 using medical records of all COVID-19 patients who had renal failure (AKI or CKD) requiring haemodialysis. The study period was from June - September 2020.

Results: Mean age of presentation was 55.08 years and with 75% of the study population were males. Most patients presented with severe COVID-19 illness with tachypnoea and hypoxia. AKI occurrence and mortality was noted in 44.1% and 47%, respectively. It was noted that medication usage was higher for corticosteroids, oseltamivir, vitamin C, zinc therapy.

Conclusion: In this single centre study involving COVID-19 patients requiring haemodialysis, it was noticed that severe COVID-19 illness, presence of AKI, chronic respiratory illness and high inflammatory markers were associated with higher mortality.

Keywords: Acute kidney injury, Cardiovascular disease, Chronic kidney disease, Diabetes mellitus

INTRODUCTION

COVID-19 presents with varied symptoms including being asymptomatic to having fever, cough, breathlessness and fatigue. Both AKI and CKD are risk factors for COVID-19 infection. COVID-19 itself can cause AKI with incidence of 3-9% and presence of AKI, CKD, hypertension, diabetes mellitus, cardiovascular disease are risk factors for poor outcome in COVID-19 infections [1,2]. Compared to normal population, renal failure patients with COVID-19 have extensive lung involvement, need more ventilator support, and have higher mortality [2]. They have mortality between 3-6% reaching almost 30% in different geographical areas [1-4].

Literature search showed that there was lack of large scale data on mortality and lack of clinical management strategies in treatment of these vulnerable patients. So, this study was expected to provide information regarding the outcome (mortality, recovery) and correlation between factors (age, gender, co-morbidities), biochemical parameters with mortality in COVID-19 patients who had renal failure (AKI or CKD) requiring haemodialysis.

MATERIALS AND METHODS

This retrospective observational study was conducted using medical records of COVID-19 patients requiring haemodialysis during period June-September 2020 at Vydehi Institute of Medical Sciences and Research Centre, Bangalore, Karnataka, India, after obtaining Institutional Ethical Committee (IEC) approval (ECR/747/Inst/KA/2015/RR-18).

Using medical records like patient's file, laboratory test reports from medical record section in the hospital, data was collected regarding patient's demography (age, gender), co-morbidities (hypertension, diabetes mellitus, pre-existent renal disease, cardiovascular disease, chronic respiratory illness), biochemical parameters like complete blood cell count, Liver function test, D-Dimer, Lactate

Dehydrogenase (LDH), ferritin, C-Reactive Protein (CRP), radiological tests like chest radiograph.

Inclusion criteria: All patients diagnosed as COVID-19 with positive pharyngeal swab Reverse Transcriptase Polymerase Chain Reaction Test (RT-PCR) who had renal failure (AKI or CKD) requiring haemodialysis.

Exclusion criteria: COVID-19 patients who had milder renal dysfunction not requiring dialysis.

COVID-19 infection was classified as mild (clinical symptoms with pneumonia), moderate (fever with pneumonia), severe (with tachypnoea with respiratory rate greater rate 30 breaths per minute, hypoxia with saturation less than 93% at rest, arterial PO₂/oxygen concentration less than 300 mmHg) [5].

Outcomes/Endpoints observed were mortality and recovery. Recovery was defined as absence of fever with normal body temperature for more than three days, significant improvement in respiratory symptoms, improved chest imaging indicating reduced inflammation and a negative RT-PCR test.

STATISTICAL ANALYSIS

The data was collected, entered and analysed with SPSS version 19.0. Continuous variables was presented as mean±Standard Deviation (SD) or median with Interquartile Ranges (IQR). Categorical variables was presented as frequency and percentage (%). Chi-square test or Fischer-exact test was performed to see the association between any two categorical variables. The p-value <0.05 was considered statistically significant. Differences in quantitative parameters in two groups were assessed by t-test.

RESULTS

There were total 68 COVID-19 positive patients requiring haemodialysis. Male population was 51 (75%) with male:female ratio of 3:1 [Table/Fig-1]. Chronic respiratory illness was noted in

40 (58.8%). Thirty (44.1%) of COVID-19 patients had AKI. Severe COVID-19 illness was noted in 27 (37.9%).

Characteristics	Survivors	Non survivors	p-value (t-test)
Number (%)	36 (53)	32 (47)	-
Mean age (years) (M±SD)	54.47±13.34	61.56±13.64	1
Gender			
Male (%)	27 (75)	24 (75)	1
Female (%)	9 (25)	8 (25)	1
Co-morbidities (%)			
Hypertension	28 (78)	25 (78)	0.97
Diabetes mellitus	20 (56)	24 (75)	0.09
Chronic respiratory illness	12 (33)	28 (87)	0.001
Cancer	1 (3)	0	0.342
Smoking	17 (47)	20 (62)	0.2
Ischemic heart disease	8 (22)	6 (19)	0.39
Old cerebrovascular accident	3 (8)	7 (22)	0.116
Presenting symptoms (%)			
Asymptomatic	2 (6)	0	0.176
Mild	17 (47)	2 (6)	0.001
Moderate	14 (39)	6 (19)	0.069
Severe	3 (8)	24 (75)	<0.001
AKI (%)	8 (22)	22 (69)	<0.001
Existent CKD (%)	22 (61)	14 (43)	0.152
Patients who skipped/ missed dialysis (%)	15 (41)	8 (25)	0.147

[Table/Fig-1]: Comparison of clinical parameters among COVID-19 survivors and non survivors (N=68).

In this study, occurrence of anaemia, high D-dimer, high LDH, hypoalbuminaemia, high serum ferritin and high CRP were noted in over more than half of the population, more frequently in mortality group [Table/Fig-2]. Further, it was noted that medication usage was higher for corticosteroids, oseltamivir, Vitamin C, Zinc therapy in mortality group [Table/Fig-2].

(a) Laboratory findings with normal cut-off values in units	Mean value among COVID-19 survivors (values in units)	Mean value among COVID-19 non survivors (values in units)	Number of patients with abnormal values among survivors n (%)	Number of patients with abnormal value among non survivors n (%)	Total number of patients with abnormal value among both groups n (%)
Anaemia (normal Hb 14-16 g/dL)	13	9.8	24 (67)	32 (100)	56 (82.4)
High D-dimer (normal 220-740 ng/mL)	367	1023	24 (67)	32 (100)	56 (82.4)
High LDH (normal <250 U/L)	261	578	24 (67)	32 (100)	56 (82.4)
High CRP (normal <10 mg/L)	9	45	25 (69)	32 (100)	57 (83.8)
Hypoalbuminaemia (normal 3.5-5 g/dL)	3.3	1.9	28 (78)	32 (100)	60 (88.2)
High S. Ferritin (normal 29-250 µg/L)	302	878	28 (78)	32 (100)	60 (88.2)
(b) Medications	Frequency among COVID-19 survivors (n, %)		Frequency among COVID-19 non survivors (n, %)		Total Percent-age among both groups (n, %)
Corticosteroids	25 (69)		32 (100)		57 (83.8)
Oseltamivir	23 (64)		32 (100)		55 (80.9)
Vitamin C	33 (92)		32 (100)		65 (95.6)
Zinc	35 (97)		32 (100)		67 (98.5)

[Table/Fig-2]: Comparison of laboratory parameters (a) and medications usage (b) among survivors (n=36) and non survivors (n=32).

When outcome was analysed, mortality and recovery were noted in 32 (47%) and 36 (53%) patients, respectively. In the mortality group [Table/Fig-1], it was noted that chronic respiratory illness, AKI, severe COVID-19 illness were seen in 28 (87%), 22 (69%), 24 (75%) patients respectively. In the survivor group [Table/Fig-1], it was noted that chronic respiratory illness, AKI, severe illness were seen in 12 (33%), 8 (22%), 3 (8%) patients, respectively. Need of Intensive Care Unit (ICU) support, and stay in hospital more than 10 days were noted in 42 (61.8%) and 29 (42.6%), respectively.

DISCUSSION

The clinical course of COVID-19 infection varies from asymptomatic to mild respiratory illness to severe disease needing ICU care, mechanical ventilation support and sometimes haemodialysis [6-10].

Co-morbidities/Risk factors: Similar to other studies, it was noted that, higher occurrence of co-morbidities like diabetes mellitus [11,12], hypertension [11,12], chronic respiratory illness [13,14] and risk factor like smoking [15] were observed in mortality group in this study.

Clinical features: In this study, higher occurrence of fatigue, fever, cough, breathlessness, both lung involvement, pneumonia was noted in mortality group like in other studies [15-18].

Complications: In this study, occurrence of AKI (44.1%) was high compared to other studies (4.5-23%) [19]. In this study, higher need of oxygen support, ICU support were noted in mortality group as such patients are prone for sepsis, shock, cerebrovascular disease, venous thromboembolism [20], multiorgan involvement including respiratory events like dyspnea, hypoxia, renal failure, altered mental status, liver dysfunction, coagulopathy, thrombocytopenia [1,20].

Treatment: In this study, it was noted that medication usage was higher for corticosteroids, oseltamivir, vitamin C, zinc therapy similar to that quoted in other studies using various medications like corticosteroids [3,19,20], vitamin C [20], oseltamivir [11,21], zinc therapy [20] [Table/Fig-3] [11,15,20,21]. Mechanism of action of these drugs quoted are direct antiviral and immunomodulatory properties [9].

Other studies	Corticosteroids	Vitamin C	Oseltamivir	Zinc therapy
Present study	57 (83.8%)	65 (95.6%)	55 (80.9%)	67 (98.5%)
Flythe JE et al., [11]	52 (36%)	10 (7%)	-	-
Rosenthal N et al., [20]	19.8%	13.6%	-	9.6%
Stefan G et al., [15]	9 (24%)	-	-	-
Ozturk S et al., [21]	12 (3.8%)	-	228 (63.7%)	-

[Table/Fig-3]: Comparison of frequency of medications used for COVID-19 among different studies [11,15,20,21].

Though several drugs, medications were under investigations during the study period, there was no effective medications nor vaccine available for COVID-19. Hence, treatment was tailored according to patient co-morbidities and clinical condition, like in other studies [15]. This shows marked heterogeneity in therapeutic approach across globe [14].

Mode of dialysis: In this study, all patients underwent single use dialyser treatment. All patients underwent regular dialysis or Sustained Low Efficiency Dialysis (SLED) according to their haemodynamic status. Continuous Renal Replacement Therapy (CRRT) was not offered due to non-affordability by patients.

Investigations: Consistent with other studies, it was noted that occurrence of anaemia, high D-dimer, high LDH, hypoalbuminaemia, high serum ferritin and high CRP were higher in mortality group [Table/Fig-4] [3,11,14,19,22]. This indicates significant role of inflammation in mortality in COVID-19.

Other studies	D-dimer	Serum albumin	Serum ferritin	CRP	LDH
Present study	1023 ng/mL	1.9 g/dL	878 µg/L	45 mg/L	578 U/L
Kooman JP and van der Sande FM [19]*	-	-	-	CRP>175 mg/dL	-
Flythe JE et al., [11]*	1347 ng/mL	-	3406 ng/mL	170 mg/L	-
Tortorese S et al., [14]*	715 IU/L	3.2 g/dL	-	65 mg/L	313 U/L
Bellan M et al., [22]*	1356 µg/mL	-	-	7.1 mg/dL	706 IU/L

[Table/Fig-4]: Laboratory findings influencing mortality in various studies [11,14,19,22].

*In all these studies, laboratory parameters were abnormal.

Outcome (Recovery and Mortality): On analysis of mortality rate, it was noted that mortality and recovery were seen in 32 (47%) and 36 (53%) patients, respectively. Interestingly it was noted that COVID-19 patients with renal failure on haemodialysis had higher mortality compared to those without [9,10,23,24]. This shows that mortality in COVID-19 was higher in presence of renal failure (AKI, CKD) [3-5,21].

In this study, mortality was related to presence of chronic respiratory illness, AKI and severe COVID-19 illness with possible underlying malnutrition and less efficient immunity in this group like in other studies [22,23,25,26].

In this study, mortality rate was 47%, higher than compared to other studies like 27.7% [23], 13.3% [10], 12% [5]. Reasons could be different baseline characters, difference in co-morbidities, study included AKI in significant numbers unlike in other studies which included predominantly CKD patients, genetic factors, variation in local testing facilities and difference in referring local health care centres.

Limitation(s)

Retrospective study, sample size was small, single centre study, follow-up time was short, and the study lacked a comparable group (COVID-19 patients without renal failure).

CONCLUSION(S)

In this single centre study, involving COVID-19 patients with renal failure requiring haemodialysis, it was noticed that mortality was associated with presence of severe COVID-19 illness, chronic respiratory illness, AKI and inflammatory biochemical markers. Further it was noticed that during this period, there was no definitive treatment for COVID-19 infection leading to random use of multiple, different medications as target therapy. In this current scenario, use of bio-chemical parameters, relating co-morbidities to determine COVID-19 prognosis needs further validation and also effective anti-viral therapy, vaccines are needed to control this COVID-19 pandemic.

REFERENCES

- Prasad N, Agrawal SK. COVID-19 and acute kidney injury. *Indian J Nephrol.* 2020;30(3):161-65.
- Ajalmy M, Melamed ML. COVID-19 in patients with kidney disease. *Clin J Am Soc Nephrol.* 2020;15(8):1087-89.
- Ikizler TA. COVID-19 in dialysis patients: Adding a few more pieces to the puzzle. *Kidney International.* 2020;98:17-19.
- Ikizler TA, Klinger AS. Minimizing the risk of COVID-19 among patients on dialysis. *Nat Rev.* 2020;16:311-13.
- Deshpande R, Dash S, Bahadur MM, Thamba A, Pathan AK, Dave K, et al. Study of COVID-19 pandemic in representative dialysis population across Mumbai, India: An observational multicentric analysis. *J Assoc Physicians India.* 2020;68(10):13-17.
- Gedney N. Long-term hemodialysis during the COVID-19 pandemic. *Clin J Am Soc Nephrol.* 2020;15:1073-74.
- Sia CSM, Cheong SHL, Ngoh CLY, Tan YH, Wong WK. Critical coronavirus disease. 2019 in a hemodialysis patient: A proposed clinical management strategy. *Case Rep Nephrol Dial.* 2020;10:86-94.
- Klinger AS, Cozzolino MC, Jha V, Harbert G, Ikizler TA. Managing the COVID-19 pandemic: International comparisons in dialysis patients. *Kidney International.* 2020;98:12-16.
- Kataria A, Yakubu I, Winstead R, Gowda M, Gupta G. COVID-19 in kidney transplantation: Epidemiology, management considerations, and the impact on kidney transplant practice. *Transplantation Direct.* 2020;1:1-10.
- Rombola G, Brunini F. COVID-19 and dialysis: Why we should be worried. *J Nephrol.* 2020.
- Flythe JE, Assimon MM, Tugman MJ, Chang EH, Gupta S, Shah J, et al. Characteristics and outcomes of individuals with pre-existing kidney disease and COVID-19 admitted to intensive care units in the United States. *Am J Kidney Dis.* 2021;77(2):190-203.
- Jager KJ, Kramer A, Chesnaye NC, Couchoud C, Sanchez-Alvarez JE, Garneata L. Results from the ERA-EDTA Registry indicate a high mortality due to COVID-19 in dialysis patients and kidney transplant recipients across Europe. *Kidney International.* 2020;98:1540-48.
- Trivedi M, Shingada A, Shah M, Khanna U, Karnik ND, Ramachandran R. Impact of COVID-19 on maintenance haemodialysis patients: The Indian scenario. *Nephrology.* 2020;25:929-32.
- Tortorese S, Scriabine I, Anjou L, Loens C, Michon A, Benabdelhak M. COVID-19 in patients on maintenance dialysis in the Paris Region. *Kidney International Reports.* 2020;5:1535-44.
- Stefan G, Mehedinti AM, Andreiana I, Zugravu AD, Cinca S, Busuioc R. Clinical features and outcome of maintenance hemodialysis patients with COVID-19 from a tertiary nephrology care center in Romania. *Renal Failure.* 2021;43(1):49-57.
- Zou R, Chen F, Chen D, Xu CL, Xiong F. Clinical characteristics and outcome of hemodialysis patients with COVID-19: A large cohort study in a single Chinese center. *Renal Failure.* 2020;42(1):950-57.
- Mazzoleni L, Ghafari C, Mestrez F, Sava R, Bivoleanu E, Delmotte P. COVID-19 outbreak in a hemodialysis center: A retrospective monocentric case series. *Canadian Journal of Kidney Health and Disease.* 2020;7:01-08.
- Mehraena E, Karimib A, Barzegaryc A, Vahedib F, Afsahid A, Dadrase O, et al. Predictors of mortality in patients with COVID-19-A systematic review. *European J Integrative Med.* 2020;40:101226.1-8.
- Kooman JP, van der Sande FM. COVID-19 in ESRD and acute kidney injury. *Blood Purif.* 2021;50(4-5):610-20.
- Rosenthal N, Cao Z, Gundrum J, Sianis J, Safo S. Risk factors associated with in-hospital mortality in a US national sample of patients with COVID-19. *JAMA Network Open.* 2020;3(12).
- Ozturk S, Turgutalp K, Arici M, Odabas AR, Altiparmak MR, Aydin Z. Mortality analysis of COVID-19 infection in chronic kidney disease, haemodialysis and renal transplant patients compared with patients without kidney disease: A nationwide analysis from Turkey. *Nephrol Dial Transplant.* 2020;35:2083-95.
- Bellan M, Patti G, Hayden E, Azzolina D, Pirisi M, Acquaviva A, et al. Fatality rate and predictors of mortality in an Italian cohort of hospitalized COVID-19 patients. *Scientific Reports.* 2020;10:20731.1-10.
- Abreu AP, Nascimento MMD, Vieira MA, Neves PDMD, Lugon JR, Sesso R. High mortality of CKD patients on hemodialysis with COVID-19 in Brazil. *Journal of Nephrology.* 2020;33:875-77.
- Jangamani R, Thirumal C, Sundar S. Coronavirus disease 2019 and the kidney. *APIK J Int Med.* 2020;8:172-78.
- Noor FM, Islam MM. Prevalence and associated risk factors of mortality among COVID-19 patients: A meta-analysis. *Journal of Community Health.* 2020;45:1270-82.
- Parohan M, Yaghoubi S, Seraji A, Hassan M, Javanbakht, Sarraf P, et al. Risk factors for mortality in patients with Coronavirus disease 2019 (COVID-19) infection: A systematic review and meta-analysis of observational studies. *The Aging Male.* 2020;01-09.

PARTICULARS OF CONTRIBUTORS:

- Associate Professor, Department of Nephrology, Vydehi Institute of Medical Sciences and Research Centre, Bangalore, Karnataka, India.
- Professor, Department of General Medicine, Vydehi Institute of Medical Sciences and Research Centre, Bangalore, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Girish P Vakrani,
'A' Block, No. 29, Vydehi Hospital Staff Quarters #82, EPIP Area Whitefield,
Bangalore-560066, Karnataka, India.
E-mail: drvakranis@gmail.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? No
- For any images presented appropriate consent has been obtained from the subjects. No

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Aug 17, 2021
- Manual Googling: Oct 12, 2021
- iThenticate Software: Oct 30, 2021 (7%)

ETYMOLOGY: Author Origin

Date of Submission: **Aug 11, 2021**
Date of Peer Review: **Sep 17, 2021**
Date of Acceptance: **Oct 28, 2021**
Date of Publishing: **Nov 01, 2021**