

# Study of Serum Sodium and Potassium in Acute Myocardial Infarction

VINOD WALI<sup>1</sup>, SINGI YATIRAJ<sup>2</sup>

## ABSTRACT

**Objectives:** Acute myocardial infarction (AMI) is one of the leading causes of morbidity and mortality across the world. Serum electrolytes changes in AMI have not been studied extensively and there is paucity of information in the literature in this regard. Scanty information is available in the literature about prognostic value of serum electrolytes in ischaemic heart disease. Hence this study was undertaken to study any changes in the serum electrolytes with special reference to serum sodium and potassium in cases of AMI and study the correlation of serum sodium and potassium in the severity and outcome of AMI.

**Materials and Methods:** Hundred people were included in study divided equally in study and control groups. Study group comprised confirmed diagnosis of recent onset of AMI. The blood samples of both the groups were analysed for Serum

electrolytes (Na<sup>+</sup>, K<sup>+</sup>) by flame-photometry (Bio-Lab Diagnostic kit).

**Results:** There was statistically significant decrease in sodium and potassium levels in across all age groups & in both sexes of study group compared to control group. Significant high level of sodium was observed in AMI patients who are smokers and AMI patients with Diabetes whereas the level was low in AMI patients with hypertension. Potassium levels were low in AMI patients with Diabetes whereas the change was insignificant in association with smoking and hypertension.

**Conclusion:** Decrease in sodium level was due to hypoxia and ischaemia, which increase the permeability of sarcolemma to sodium whereas decrease in potassium level was influenced by the catecholamine levels which are elevated in early acute myocardial infarction.

**Keywords:** Electrolytes, MI, Sodium, Potassium

## INTRODUCTION

Cardiovascular disease is one of the leading causes of morbidity and mortality across the world. World Health Organization (WHO) has declared cardiovascular disease as a modern epidemic. AMI is one of the manifestations of coronary heart disease leading to morbidity and mortality. In India, the prevalence of ischemic heart disease among adults (based on clinical and ECG criteria) was estimated at 96.7 per 1000 population in the urban and 27.1 percent in rural areas [1].

Several systemic metabolic changes occur in AMI [2]. These changes include increased plasma concentrations of catecholamines, free fatty acids, glucose, glycerol, cortisol and cyclic-AMP. There is decreased triglycerides concentration and an initial fall in plasma insulin concentration, followed by an early return to normal value [3].

Serum electrolytes changes in AMI have not been studied extensively and there is paucity of information in the literature in this regard. Scanty information is available in the literature about prognostic value of serum electrolytes in ischaemic heart disease.

## AIMS AND OBJECTIVES

The study was undertaken to detect any changes in the serum electrolytes with special reference to serum sodium and potassium in cases of AMI and study the correlation of serum sodium and potassium in the severity and outcome of AMI.

## MATERIALS AND METHODS

After obtaining the clearance from Ethical Committee of the Institution, the study was carried out on 50 patients with recent onset of acute myocardial infarction according to European Society of Cardiology (ESC), the American College of Cardiology (ACC), the American Heart Association (AHA) and the World Heart Federation (WHF) jointly [4], admitted in the Department of Medicine, during

the period May 2008 to October 2009 at Basaveshwar Teaching & General Hospital attached to M R Medical College Gulbarga, Karnataka, India.

### Study group

The patients in study group were selected based on age, sex, duration of symptoms of AMI and any history suggestive of vascular complications with the help of physician's opinion.

### Control group

For the control purpose, 50 normal non-hypertensive persons without symptoms of AMI were carefully selected and examined in detail with age and sex matched, who were not obese, non predisposed and physically active.

### Informed consent

Informed consent was taken from both study and control group subjects after explaining the purpose and procedure of the study.

### Collection of samples

Collection of venous blood samples was done in both the groups on the day of admission within 12 hours from antecubital vein with all aseptic precautions in plain vacutainers for serum electrolytes, i.e., Na<sup>+</sup> & K<sup>+</sup>.

Blood was allowed to clot at room temperature for half an hour and then centrifuged at 3000 rpm for five minutes. The serum separated was used for the estimation.

Serum electrolytes (Na<sup>+</sup>, K<sup>+</sup>) were measured by flame-photometry (Bio-Lab Diagnostic kit).

The comparative study of serum and potassium levels was carried out between the controls and AMI, patients and AMI patients with and without history of Smoking, Hypertension and Diabetes

Sr. No.	Parameter	<50 years				>50 years			
		Control (n= 18)	AMI (n= 14)	z-value	p-value	Control (n=32)	AMI (n= 36)	z-value	p-value
1	Sodium (mEq / L)	137.88 ±3.46	130.71 ± 5.71	4.14	<0.001*	136.53 ± 3.42	129.47 ± 4.87	6.993	<0.001*
2	Potassium (mEq / L)	4.10 ± 0.81	3.31 ± 1.12	2.32	< 0.01*	4.67 ± 0.90	4.18 ± 0.63	2.684	< 0.01*

[Table/Fig-1]: Serum sodium and potassium levels in age groups (<50 years and >50 years) of AMI and normal healthy subjects, \* = shows 'Z' test is statistically significant

Sr. No.	Parameter	<50 years				Female			
		Control (n= 27)	AMI (n= 28)	z-value	p-value	Control (n=23)	AMI (n= 22)	z-value	z-value
1	Sodium (mEq / L)	137.22 ± 3.53	129.96 ± 5.84	5.606	<0.001*	136.78 ± 3.45	129.63 ± 4.07	6.349	<0.001*
2	Potassium (mEq / L)	4.18 ± 0.75	3.75 ± 0.92	8.431	<0.001*	4.13 ± 0.58	3.28 ± 1.03	13.531	<0.001*

[Table/Fig-2]: Serum sodium and potassium levels in different sex groups of AMI and normal healthy subjects, \* = shows 'Z' test is statistically significant

Sr. No.	Parameter	AMI without smoker (n=40)	AMI with smoker (n=10)	z-value	p-value
1	Sodium (mEq / L)	128.40 ± 4.60	135.5 ± 2.10	7.244	p < 0.001*
2	Potassium (mEq / L)	4.19 ± 0.67	4.06 ± 0.68	0.543	p > 0.05**

[Table/Fig-3]: Serum sodium and potassium levels in AMI with smokers and AMI with non-smokers of subjects, \* = shows 'Z' test is statistically significant ; \*\* = Shows 'Z' test is statistically non-significant

Sr. No.	Parameter	AMI without smoker (n=23)	AMI with smoker (n=27)	Z value	P value
1	Sodium (mEq / L)	130.13 ± 5.69	129.55 ± 4.62	0.391	P > 0.05**
2	Potassium (mEq / L)	4.01 ± 0.62	4.29 ± 0.71	0.540	P > 0.05**

[Table/Fig-4]: Serum sodium and potassium levels in AMI with hypertension and AMI without hypertension of subjects., \*\* = shows 'Z' test is statistically non-significant

Sr. No.	Parameter	AMI without smoker (n=39)	AMI with smoker (n=11)	z-value	p-value
1	Sodium (mEq / L)	129.51 ± 5.03	130.90 ± 4.32	0.90	p > 0.05**
2	Potassium (mEq / L)	4.17 ± 0.73	3.69 ± 0.90	1.632	p > 0.05**

[Table/Fig-5]: Serum sodium and potassium levels in AMI with diabetes mellitus and AMI without diabetes mellitus of subjects, \*\* = shows 'Z' test is statistically non-significant

Mellitus, also a study was carried out to evaluate the variations in serum electrolyte levels in AMI patients of age below 50 years and above 50 years.

**Inclusion criteria**

- a) Patients of either sex and of any age group,
- b) Patients diagnosed clinically by Physician based on Clinical examination and ECG changes.

**Exclusion criteria**

- a) Patients on diuretics,
- b) Patient not fitting into criteria.

**STATISTICAL ANALYSIS**

The data of 50 AMI patients and 50 non-AMI persons (Control) thus collected were analysed, correlation of serum Na<sup>+</sup> and K<sup>+</sup> parameters expressed as cross-tabulations. The mean and standard deviations of SBP, DBP serum electrolytes (Na<sup>+</sup>, K<sup>+</sup>) were calculated and the 'cut-off' points taken for finding a normal range. Measurement data in the form of cross-tabulations for observed values were found out and the simple 'Z' test standard error of difference between two proportions used as tests of significance.

**RESULTS**

The AMI cases were observed in Intensive Cardiac Care Unit and their serum electrolytes value were estimated and these values were compared with normal healthy control group.

There was statistically significant decreased in serum sodium and potassium levels in study group across both all ages compared to normal healthy control group. A significant low level of potassium was observed in <50 years of age group patients of AMI than normal healthy controls, but there was higher level of potassium observed in > 50 years age groups of AMI cases [Table/Fig-1]. This statistically significant decrease in serum electrolyte levels was also seen in both sexes [Table/Fig-2].

Significantly high level of serum sodium was observed in AMI with smokers than AMI with non-smokers. No significant difference was found in the level of serum potassium in AMI with smokers and non-smokers [Table/Fig-3]. Whereas levels of serum sodium was decreased in AMI with hypertension than AMI without hypertension. No significant change was observed in serum potassium level in AMI with hypertension and AMI without hypertension [Table/Fig-4]. However this comparison was statistically insignificant. Similar statistically insignificant findings were noted in AMI with diabetes mellitus and AMI without diabetes mellitus. Serum sodium levels were higher in AMI with diabetes mellitus than AMI without diabetes mellitus. Lower value of serum potassium was observed in AMI with diabetes mellitus as compared to AMI without diabetes mellitus [Table/Fig-5].

**DISCUSSION**

Fall in sodium level on day of admission in AMI is similar to finding was observed by Shah and his coworkers et al., [5] who reported hyponatremia on day first. Flear and Hilton reported a progressive fall in the mean daily serum sodium concentration until day 4 and rise thereafter in all cases [6]. Significant decrease in serum sodium concentration in both sexes was also reported by Flear and Singh [7].

There were 10 cases who were smokers in which the serum sodium concentration was significantly higher than non-smokers. No support is available in literature to show the correlation of smoking and sodium concentration.

The low level of serum potassium was observed in below 50 years of age groups patients than above 50 years. There was a tendency for potassium levels to increase with age [8]. Low level of serum potassium was seen in AMI cases in both the age groups. Flear and Hilton reported fall in mean daily serum potassium concentrations during first three days of AMI [6].

The level of serum potassium in female cases was significantly decreased than males. The higher frequency of hypokalemia in women than in men in AMI has been reported by others, but the cause has not been known [9-10].

Serum potassium level was non-significantly higher in non-smokers than smokers. Smoking showed a strong association with serum potassium level. Higher level was reported by Wannamethee and his coworkers in normal middle aged person [8]. They also reported increase in mortality with increasing serum potassium levels in cardiovascular disease.

The level of serum potassium was non-significantly increased in AMI with hypertension than AMI without hypertension cases. Hypokalemia is not a feature of hypertension but the plasma potassium level is invariably increased in hypertensive patients [11]. Hamid Ikram and his colleagues reported low level of serum potassium level in hypertensive coronary heart disease [12]. Decrease level of serum potassium was observed in AMI with diabetes mellitus, which is non-significant.

## CONCLUSION

Hyponatremia and hypokalemia are indicators of Acute myocardial infarction. Serum sodium and potassium levels are prognostic indicators, i.e, rise in sodium levels after initial fall was indicative of clinical improvement. Therefore estimation of sodium and potassium level in acute MI patients can help assess their prognosis.

## REFERENCES

- [1] Park. Textbook of Preventive and Social Medicine 19<sup>th</sup> Ed. 2007: 286-287.
- [2] Oliver MP. Metabolic response during impending myocardial infarction, II Clinical implications. *Circulation*. 1972; 45: 491.
- [3] Vetter NJ, Adams W. Strange RC, Oliver MF: Initial metabolic and hormonal response to AMI. *Lancet*. 1974; I: 284.
- [4] Thygesen K, Alpert JS, White HD, Jaffe AS, Apple FS, Galvani M, et al. Universal definition of myocardial infarction Kristian Thygesen, Joseph S. Alpert and Harvey D. White on behalf of the Joint ESC/ACCF/AHA/WHF Task Force for the Redefinition of Myocardial In-farction. *Eur Heart J*. 2007 Jan 10; 28(20):2525-38.
- [5] Shah PKD, Saxena M, Khanqarot D, Banerji K.: Study of serum sodium in myocardial infarction and its relation to severity and complications. *JPI J Assoc Physicians India*. 1986;34(3):195.
- [6] Flear CTG, Hilton P.: Hyponatremia and severity and outcome of myocardial infarction. *Br. Med. J*. 1979; I:1242.
- [7] Flear CTG, Singh CM. Hyponatremia and sick cells. *Br. J. Anaesth*. 1973;45: 976.
- [8] Wannamethee SG, Lever AF, Shaper AG, Whincup PH. Serum potassium, cigarette smoking and mortality in middle aged men. *Am. J. of Epidemiology*. 1997;145(7):598.
- [9] Dyckner T, Helmers C, Lundman T, Wester PO. Initial potassium level in relation to early complications and prognosis in patients with AMI. *Acta Med. Scand*. 1975;197:207-10.
- [10] Bech OA, Hochrein H. Serumkaliumspiegel und Herzrhythmusstorungen beim akuten Myokardinfarkt Z. *Kardiol*. 1977;66:187.
- [11] Walmsley RN, Guerin MD: Disorders of fluid and electrolyte balance. 1984: WRIGHT; 244.
- [12] Ikram H, Espiner EA, Nicholls MG. Diuretics, potassium and arrhythmias in hypertensive coronary disease. *Drug*. 31(Suppl 4);101.

### PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Biochemistry, M. R. Medical College, Gulbarga, Karnataka, India.
2. Associate Professor, Department of Forensic Medicine, Sree Mookambika Institute of Medical Sciences, Kulasekharam, TN, India.

### NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Vinod Wali,  
Assistant Professor, Department of Biochemistry, M. R. Medical College; Sedam Road, Gulbarga-585101, Karnataka, India.  
E-mail : docvinod80@yahoo.com

FINANCIAL OR OTHER COMPETING INTERESTS: None.

Date of Submission: **Jun 23, 2014**  
Date of Peer Review: **Aug 21, 2014**  
Date of Acceptance: **Oct 01, 2014**  
Date of Publishing: **Nov 20, 2014**