

To Study the Correlation of Thompson Scoring in Predicting Early Neonatal Outcome in Post Asphyxiated Term Neonates

DALIP KUMAR BHAGWANI¹, MANISHA SHARMA², STANZIN DOLKER³, SHARADA KOTHAPALLI⁴

ABSTRACT

Introduction: Throughout the world each year, an estimated 23% of the 4 million neonatal deaths and 8% of all deaths in <5 years of age are associated with signs of asphyxia at birth.

Aim: To study the role of cord arterial blood gas analysis at birth and serial Thompson score (HIE score) in predicting the early neonatal outcome in post asphyxiated term neonates.

Materials and Methods: The study was conducted in Department of Paediatrics, in Neonatal Intensive Care Unit (NICU), Hindu Rao Hospital, New Delhi from May 2014 to February, 2015. This study was a prospective cross-sectional study. During this period, a total of 145 post asphyxiated term neonates born in labour room/obstetric operation theatre were recruited. An informed consent was taken from all the parents. The protocol was approved by the institutional ethical committee. Inclusion Criteria were Full-term babies with low-Apgar score i.e., 1 min score of ≤ 7 (NNPD2010).

Statistical Analysis: SPSS 17.0 Software has been used for data analysis. The data were expressed in terms of Means, Standard Deviation and Proportion, followed by comparison between groups through chi-square test or Fisher's-exact test. A p-value of less than 0.05 was considered as statistically significant.

Results: The present study was carried out on 145 post asphyxiated full-term babies with low-Apgar score i.e., 1min score of ≤ 7 mild Thompson score on day 1,2,3 were 96 (66.2%), 119 (82.06%), 125 (86.20%), moderate Thompson score on day 1,3, 7 were 13 (8.9%), 6 (4.13%), 2 (1.37%) and severe Thompson score on day 1, 3, 7 were 36 (24.8%), 13 (8.96%), 7 (4.82%) respectively. Total 11 patients died out of 145 post asphyxiated full-term babies within 7 days, among 11 patients, 7 died within 3 days. There was clinical improvement among HIE patients as indicated by serial Thompson score done on day 1, 3 and 7. Among 145 patients 62(42.8%) had seizure and 83(57.2%) did not have seizure. Most common type of seizure was subtle seizure in 25 (40.3%) followed by multifocal in 21 (33.9%) and tonic in 16(25.8%).

Conclusion: There is statistically significant correlation between morbidity and day 1 Thompson score (p-value 0.024). There is statistically significant correlation between mortality and day 1 Thompson score (p-value 0.001).

Thompson score allows a very precise description of infants by assigning a numeric score rather than 'mild', 'moderate' or 'severe'. Inter-rater reliability is very good with a kappa coefficient of 0.87.

Keywords: Cord blood gas, Hypoxic Ischemic Encephalopathy, Maternal risk factor, Seizures

INTRODUCTION

Birth asphyxia is defined as a delay in establishing spontaneous breathing/crying immediately after birth of the newborn [1]. This results in impaired gas exchange, hypoxia and metabolic acidosis [2]. But it is probably better to use the term perinatal asphyxia since asphyxia may occur in utero, at birth or in the postnatal period. Hypoxic Ischemic Encephalopathy (HIE) is the term commonly used to describe the neurological syndrome that occur following perinatal asphyxia. It is usually caused by severe birth asphyxia with secondary cerebral ischemia [3]. It is an important cause of permanent damage to the Central Nervous System (CNS) which may result in neonatal death or manifest later as cerebral palsy or mental deficit [4]. Throughout the world each year, an estimated 23% of the 4 million neonatal deaths and 8% of all deaths in <5 years of age are associated with signs of asphyxia at birth [5]. Most of these infants are born in developing nations, and more than 50% of the deaths occur at home, where majority of these infants are born [6]. A 15%-20% of HIE cases die during the neonatal period and 30% of those who survive, suffer from neurodevelopment disorders [5,7,8]. In less developed countries perinatal asphyxia remains a major cause of death and disability.

The pattern of risk factors, the nature of sequelae and the options and priorities for intervention (both preventive and therapeutic) are significantly different than in the industrialized countries [9]. The HIE score (Thompson score) is a clinical tool comprising of a set of clinical signs associated with CNS dysfunction. It is used to assess status of a child following birth asphyxia [10-13]. In the scoring system, a score of 0 is normal and the maximum score is 22 which signifies the worst possible status of HIE. Infants with score 1-10 are considered to have mild HIE, 11-14 have moderate HIE and 15-22 are considered to have severe HIE. In normothermic infants, a maximum score of > 10 during the first 7 days of life predicts an abnormal outcome with 100% sensitivity and 61% specificity [14]. This study was done to assess the role of cord arterial blood gas analysis at birth and serial Thompson score (HIE score) in predicting the early neonatal outcome in post asphyxiated term neonates.

MATERIALS AND METHODS

The study was conducted in Department of Paediatrics, in NICU, Hindu Rao Hospital, New Delhi from May 2014 to February, 2015. This study was a Prospective cross-sectional study. During

this period, a total of 145 post asphyxiated term neonates born in labour room/obstetric operation theatre were recruited. An informed consent was taken from all the parents. The protocol was approved by the institutional ethical committee.

Inclusion Criteria: Full-term babies with low-Apgar score i.e., 1 min score of ≤ 7 . (NNPD) [15].

Exclusion Criteria: Preterm babies, Respiratory depression due to intracranial bleed, Neonates with major congenital malformation of CVS, CNS, Respiratory system or dysmorphic babies, Severe hyperbilirubinemia bordering on kernicterus, Case with hypoglycaemia or meningitis as cause of encephalopathy.

Criteria to define HIE: HIE characterized by Thompson score above 5 one hour after birth [16]. The score is derived from nine aspects of the neurological examination of infants with HIE, shown in [Table/Fig-2].

Cord arterial blood gas analysis was done at birth (pH, pCO₂, Bicarbonate, base deficit). Arterial cord blood samples were collected from double clamped umbilical cord, anaerobically, using heparinized disposable syringes (2ml syringe washed by 1000IU/ml heparin) and the samples were analysed by Blood Gas Analyser –Bayer, Germany. It directly measures pH, pCO₂, bicarbonate, base deficit. All infants with severe post-asphyxial HIE had evidence of dysfunction of at least one organ/system in addition to the central nervous system. This conforms with the criteria of the American College of Obstetricians and Gynaecologists [17] and Society of Obstetricians and Gynaecologists of Canada [18].

Criteria to define outcome: Outcome measures were grouped as [19]. Normal, Morbidity, neonatal encephalopathy with seizures, neonatal encephalopathy with seizures requiring anticonvulsant medication, neonatal encephalopathy with seizures requiring medication, within 24 hours of birth, apgar score < 7 at 5 minutes and admission to the neonatal unit and death within 4 weeks.


STATISTICAL ANALYSIS

SPSS 17.0 software has been used for data analysis. The data were expressed in terms of Means, Standard Deviation and Proportion, followed by comparison between groups through chi-square test or Fisher' p-value of less than 0.05 was considered as statistically significant.

RESULTS

Apgar score of ≤ 3 at 1minute was found in 58 babies (40%) and 87 babies (60%) had a score of 4-6 at 1 minute.

Mild Thompson score on day 1,3,7 were, 96 (66.2%), 119 (82.06%), 125 (86.20%), moderate Thompson score on day 1,3, 7 were 13 (8.9%), 6 (4.13%), 2 (1.37%) and severe Thompson score on day 1, 3, 7 were 36 (24.8%), 13 (8.96%), 7 (4.82%) respectively as given in [Table/Fig-3]. Total 11 patients died within 7 days among



[Table/Fig-1]: Selection of data by inclusion and exclusion criteria.

Score Sign	0	1	2	3
Tone	Normal	Hyper	Hypo	Flaccid
LOC	Normal	Hyperalert, stare	Lethargic	Comatose
Fits	None	$< 3/day$	$> 2/day$	
Posture	Normal	Fisting, cycling	Strong distal flexion	Decerebrate
Moro	Normal	Partial	Absent	
Grasp	Normal	Poor	Absent	
Suck	Normal	Poor	Absent \pm bites	
Respiration	Normal	Hyperventilation	Brief apnea	IPPV (apnea)
Fontanelle	Normal	Full, not tense	Tense	

[Table/Fig-2]: The thompson HIE score [14].

Thompson score	Day 1	Day 3	Day 7
	n = 145	n = 145	n = 145
Mild (1 - 10)	96	119	125
Moderate (11 - 14)	13	6	2
Severe (15+)	36	13	7

[Table/Fig-3]: Grading and evolution of thompson score at Day 1, 3 & 7 (improvement/survival rate).

Apgar score at 5 min	Cord gas acidosis			P
	Yes	No	Total	
	n = 145 (%)	n = 145 (%)	n = 145 (%)	
≤ 3	16 (29.6)	1 (1.1)	17 (11.7)	< 0.001
> 3	38 (70.4)	90 (98.9)	128 (88.3)	
Total	54(100)	91(100)	145(100)	

[Table/Fig-4]: Correlation between 5 minute Apgar score and cord gas acidosis.

Seizures	HIE				P
	Mild	Moderate	Severe	Total	
	n = 145 (%)	n = 145 (%)	n = 145 (%)	n = 145 (%)	
No	67(69.8)	1(7.7)	15(41.7)	83(57.2)	< 0.001
Yes	29(30.2)	12(92.3)	21(58.3)	62(42.8)	
Total	96(100)	13(100)	36(100)	145(100)	

[Table/Fig-5]: Co-relation between HIE score on day 1 and neonatal convulsion.


Types of seizures	Subtle	Tonic	Multifocal	Total
	n = 62 (%)	n = 62 (%)	n = 62 (%)	
No. (%)	25(40.3%)	16(25.8%)	21(39.9%)	62(100%)

[Table/Fig-6]: Co-relation between HIE score on day 1 and neonatal convulsion.


	Phenobarbitone loading dose	+ Pheno-barbitone maintenance dose	+ phenytoin Loading dose	+ phenytoin Maintenance Dose	+ Midazolam infusion	Total
	n = 62 (%)	n = 62 (%)	n = 62 (%)	n = 62 (%)	n = 62 (%)	
No. of babies	50 (80.6)	5 (8%)	3 (4.83)	2 (3.22)	2 (3.22)	62

[Table/Fig-7]: Babies who required anticonvulsants.

them 7 patients died within 3 days. There was clinical improvement among HIE patients as indicated by serial Thompson score done on day 1, 3 and 7. Out of 145 patients, cord gas acidosis was found in 54 patients as given in [Table/Fig-4], there is statistically significant correlation between morbidity and day 1 Thompson score (p-value 0.024). There is statistically significant correlation between mortality and day 1 Thompson score (p-value 0.001). There is a statistically significant relation between duration of



[Table/Fig-8]: ROC curve (Morbidity with Thompson score).



[Table/Fig-9]: ROC curve (Mortality with Thompson score).

hospital stay and HIE grading ($p=0.008$) with maximum duration of stay in severe HIE cases (mean value 7.64 days) total 62 (42.8%) patients had seizures. The type of seizures and management is as given in [Table/Fig-5-7].

In 145 cases as depicted in [Table/Fig-8,9], ROC curves were plotted for Thompson score day 1 and morbidity and Thompson score day1 and mortality. For the mortality curve, AUC (Area under Curve) was 0.936 which was highly significant ($p=0.001$). There was a tendency for better predictive capacity of Thompson score in regard to morality (Sensitivity 93%, Specificity 90%, Positive Predictive Value (PPV) 50% and Positive Predictive Value (NPV) 99%). For the morbidity curve, AUC (Area under Curve) was 0.561 which was not significant ($p=0.324$), Sensitivity was 63%, Specificity was 40%, NPV was 19% and Negative Predictive Value was 82%.

DISCUSSION

The present study was carried out on 145 newborns with perinatal asphyxia admitted to NICU, department of pediatrics Hindu Rao Hospital Delhi. For delivery room resuscitation, the standard NRP AAP 2010 guidelines were followed and cord arterial blood gas analysis was done in each patient. The cases were further divided into mild, moderate and severe HIE, according to Thompson score.

In present study 57 (39.3%) patients had maternal risk factor. The most common maternal risk factor was those mother who didn't

register ANC (Anti Natal Checkup) as given for in [Table/Fig-10] [20-22].

MSL (Meconium Stained Liquor) was the most common intrapartum risk factor seen in 54 (37.2%) cases which was in accordance with Dongol S et al., [20] (37.2%), but higher than Padayachee N et al., [23] (10.9%) and lesser than Shrestha M et al., [24] (65%). Post term as an intrapartum risk factor was present in 10 (6.9%). Our finding is similar to Mundhra R et al., [21] (6.06%), Qureshi AM et al., [25] (6%). We found prolonged labour in 8 (5.5%) which was lesser than Qureshi AM et al., [25] (9.3%).

Maternal risk factor	Maternal age ≥ 35 yrs	No/<3 Antenatal Visit	PIH (Pregnancy induced hypertension)	BOH (Bad Obstetric History)	Gestational Diabetes	Total	No maternal risk factor
No. of babies n=145 (%)	15 (10.3)	29 (20)	16 (11)	8 (5.5)	7 (4.8)	57 (39.3)	88 (60.7)
Author	Dongol S et al., [20] (9.8%), Mundhra R et al., [21] (23.6%)	Dongol S et al., [20] (15.68%)	Mundhra R et al., [21] (16.97%), Kerkhofs C et al., [22] (9.4%).	-----	Kerkhofs C et al., [22] (5.7%).	-----	

[Table/Fig-10]: Comparative study of Relationship between maternal risk factor and perinatal Asphyxia [20-22].

Study	Findings
Malin et al., [26]	Low arterial pH was strongly associated with neonatal mortality and long-term adverse outcomes.
Haddad et al., [27]	Arterial cord pH <7 was associated HIE and death within neonatal period.
Ghosh et al., [28]	Arterial cord pH ≤ 7.15 was associated with Hypoxic ischaemic encephalopathy and death during neonatal period.
Graham EM et al., [29]	109 33 (30.3%) HIE 27 (24.8%) seizures 5 (4.6%) deaths
Present study	54 17.7% had mild HIE, 53.8% had moderate HIE and 83.3% had severe HIE

[Table/Fig-11]: Association of Cord gas acidosis and HIE in prediction of outcomes [26-29].

There is strong association of cord gas acidosis (pH <7) and severity of HIE in the present study as given in [Table/Fig-11] [26-29]. In normothermic infants, Thompson score of >10 during the first 7 days of life predicts an abnormal outcome with 100% sensitivity and 61% specificity [14].

LIMITATION

Longterm neurodevelopmental outcome was not assessed in our study and outcome prediction could better be assessed with the help of neuroimaging.

CONCLUSION

Thompson score allows a very precise description of infants by assigning a numeric score rather than 'mild', 'moderate' or 'severe'. Inter-rater reliability is very good with a kappa coefficient of 0.8 and there is no requirement of Electroencephalogram (EEG) which is beneficial in resource limited country like ours. Acid-base analysis in the first hour of life suggests that a threshold level of the pH and base deficit may indicate the presence of intrapartum hypoxia more specifically and objectively than a low Apgar score or an abnormal CTG.

REFERENCES

- [1] Airede AI, Weerasinghe HD. Birth asphyxia: a review. *East Afr Med J*. 1995;72:252-57.
- [2] JA L. Intrapartum fetal asphyxia; definition, diagnosis and classification. *AM J Bbstet Gynaecol*. 1997;176:957-59.
- [3] Coovadia HM WD. Paediatrics and Child Health: A Manual for Health professionals in the third world: 4th Ed 1998:129-31.
- [4] Sperling M. Hypoxia –ischemia. Nelson's Text book-71. of Paediatric.
- [5] Stoll BJ, Kliegman RM. Hypoxia ischemia. Nelson Textbook of Pediatrics. 18th ed. Philadelphia; Saunders. 2007; Pp:566-68.
- [6] Haider BA, Bhutta ZA. Birth Asphyxia in Developing Countries: Current Status and Public Health Implications. *Curr Probl Pediatr Adolesc Health Care*. 2006;36:178-88.
- [7] Eghbalian F, Monsef A. Neonatal seizures; Etiology and frequency. *Iran J Child Neurol*. 2008;2(1):39-42.
- [8] Sghbalian F. Diagnosis natal seizures and. 1st managemented, Hamadan University of Medical neo Sciences. 2004; Pp 60-76.
- [9] Costello AM, Manandhar DS. Perinatal asphyxia in less developed countries. *Arch Dis Child Fetal Neonatal*. Ed 1994; F1-F3.
- [10] Sarnat HB, Sarnat SM. Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. *Arch Neurol* 1976;83:696-705.
- [11] Thompson CM, Puterman AS, Linley LL, Hann FM, van der Elst CW, Molteno CD, et al. The value of the Hypoxic Ischaemic Encephalopathy score in predicting neurodevelopmental outcome. *Acta Paediatrica*. 1997;86:757-61.
- [12] Lipper EG, Voorhies TM, Ross G, Vannucci RC, Auld PA.. Early predictors of one Year outcome for infants asphyxiated at birth. *Dev Med Child Neurol Suppl*. 1986;28:303-09.
- [13] Boo NY, Chandran V, Zulfiqar MA, Zamratol SM, Nyein MK, Haliza MS, et al. Early cranial ultrasound changes as predictors of outcome during first year of life in term infants with perinatal asphyxia. *J Paediatr Child Health*. 2000;36:363-69.
- [14] Thompson CM, Puterman AS, Linley LL, Hann FM, van der Elst CW, Molteno CD, et al. The value of a scoring system for hypoxic ischaemic encephalopathy in predicting neurodevelopmental outcome. *Acta Paediatrica*. 1997;86:757-61.
- [15] Report of the National Neonatal Perinatal Database 2010 Newbornwhocc.org.
- [16] Biselele T, Nau laers G, Tady B. Evolution of the Thompson score during the first 6 h in infants with perinatal asphyxia. *Acta Paediatr*. 2014;103:145-48.
- [17] ACOG Committee Opinion. Inappropriate uses of the terms fetal distress and birth asphyxia. *Int J Gynaecol Obstet*. 1998;61:309-10.
- [18] Policy statement: Task force on cerebral palsy and neonatal asphyxia (part 1). *J SOGC*. 1996;18:1267-79.
- [19] Yeh P, Emary K, Impey L. The relationship between umbilical cord arterial pH and serious adverse neonatal outcome: analysis of 51,519 consecutive validated samples. *BJOG*. 2012;119:824-31.
- [20] Dongol SSJ, Shrestha S, Shakya A. Clinical profile of birth asphyxia in dhulikhel hospital: a retrospective study. *J Nepal Paediatr Soc*. 2010;30:141-46.
- [21] Mundhra R, Aggarwal M. Fetal Outcome in Meconium Stained Deliveries. *Journal of Clinical and Diagnostic Research*. 2013;7:2874-76.
- [22] Kerkhofs C, De Bruyn C, Mesens T, Theyskens C, Vanhoestenbergh M, Bruneel E, et al. Identification of peripartum near-miss for perinatal audit. *Facts Views Vis Obgyn*. 2014;6:177-83.
- [23] Padayachee N, Ballot DE. Outcomes of neonates with perinatal asphyxia at a tertiary academic hospital in Johannesburg, South Africa. *SAJCH*. 2013;7:89-94.
- [24] Shrestha M, Shrestha L, Shrestha PS. Profile of asphyxiated babies at Tribhuvan University Teaching hospital. *J Nep Paediatr Soc*. 2009;29(1):3-5.
- [25] Qureshi AM, Ur Rehman A, Siddiqi TS. Hypoxic ischemic encephalopathy in neonates. *J Ayub Med Coll Abbottabad*. 2010;22:190-93.
- [26] Malin GL, Morris RK, Khan KS. Strength of association between umbilical cord pH and perinatal and long term outcomes: systematic review and meta-analysis. *BMJ*. 2010;340:c1471.
- [27] Haddad B, Mercer BM, Livingston JC, Talati A, Sibai BM. Outcome after successful resuscitation of babies born with apgar scores of 0 at both 1 and 5 minutes. *Am J Obstet Gynaecol*. 2000;182:1210-14.
- [28] Ghosh BM. Prediction of perinatal asphyxia with nucleated red blood cells in cord blood of newborns. *Int J Gynaecol Obstet*. 2003;81:267-71.
- [29] Graham EM, Ho lcroft CJ, Rai KK, Donohue PK, Allen MC. Neonatal cerebral white matter injury in preterm infants is associated with culture positive infections and only rarely with metabolic acidosis. *Am J Obstet Gynaecol*. 2004;191:1305-10.

PARTICULARS OF CONTRIBUTORS:

1. Senior Pediatrician, Department of Paediatrics, Hindu Rao Hospital, Delhi, India.
2. Senior Specialist, Department of Obstetrics and Gynaecology, Hindu Rao Hospital, Delhi, India.
3. Senior Resident, Department of Paediatrics, Hindu Rao Hospital, Delhi, India.
4. CMO(SAG), Department of Paediatrics, Hindu Rao Hospital, Delhi, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Dalip kumar Bhagwani,
Senior Pediatrician, Department of Paediatrics, Hindu Rao Hospital, Delhi -110085, India.
E-mail: drdalip@yahoo.com

FINANCIAL OR OTHER COMPETING INTERESTS: None.

Date of Submission: **Jul 20, 2016**
Date of Peer Review: **Aug 20, 2016**
Date of Acceptance: **Sep 19, 2016**
Date of Publishing: **Nov 01, 2016**