

Cardiac Myeloid Sarcoma: Review of Literature

ARCHANA GAUTAM¹, GHAZAL KOOSHK JALALI², KAMAL KANT SAHU³, PRATEEK DEO⁴, SIKANDER AILAWADHI⁵

ABSTRACT

Granulocytic Sarcomas (GS) also called as Myeloid Sarcomas (MS) or chloromas are the representatives of extramedullary infiltrates of immature myeloid cells including myeloblasts, promyelocytes and myelocytes. Primary cardiac malignancies per se are rare and infiltration of cardiac muscles by secondary malignant cells is also an uncommon finding. Out of these cardiac tumors, contribution of Cardiac Myeloid Sarcoma (CMS) is even more smaller thereby limiting our knowledge about this rare entity. Because of its very lower incidence, an exact guideline for diagnosis and management is still missing and usually haematologists around the world are treating CMS based on their clinical acumen. Aim of this review is to briefly discuss the presenting clinical feature, differential diagnosis, diagnostic workup and management based on published articles related to CMS till date.

Keywords: Acute myeloid leukaemia, Granulocytic sarcoma, Chemotherapy

INTRODUCTION

GS is most commonly associated with Acute Myeloid Leukaemias (AML), but can be also seen in association with Myeloproliferative Neoplasms (MPN) or Myelodysplastic Syndromes (MDS) [1]. Very rarely, it has been reported in chronic myelogenous leukaemia presenting in blast crisis. The usual reported location of chloroma is skin, bone, periosteum, spleen and soft tissues. However, acute leukaemias especially AML can have atypical presentations and we have reported rare areas of involvement such as orbit, pleura, mediastinum and spinal cord in recent past [2-7]. Presence of CMS either as locally invasive or isolated cardiac intracavitary mass is highly uncommon. The occurrence has been reported anytime during the entire disease course i.e., before the diagnosis, during treatment, post chemotherapy relapse or post stem cell transplant [8]. In most cases, symptoms are similar to that of classical commoner cardiac illnesses [9]. Both the diagnosis as well as management is challenging and must be tailored based on clinician's experience and patient's performance status. MS in other common locations is well known findings in AML and treatment protocols have also been laid down. In view of its rarity of occurrence, diagnostic limitations and restricted use of conventional anti-leukaemia agents (due to the fear of cardio toxicity), the same guidelines cannot be applied to CMS. This review aims to highlight the challenges which haematologists do face while dealing with such cases.

MATERIALS AND METHODS

We did a MEDLINE/PubMed search with MeSH terms—"Heart" and "Myeloid Sarcoma" and "(Cardiac and Myeloid Sarcoma)". In total, we found 36 articles reported (From 1960 to July 2016). For a comprehensive review, we have included all the cases including both paediatric and adult population. Exclusion criteria was removal of reports from review which were written in languages other than English. Hence, six cases (written in Japanese, French, Italian, and German language) were excluded. Hence, finally a total number of 30 articles were included in this study [Table/Fig-1]. Detailed summary of all the articles included in the study is mentioned in [Table/Fig-2] [1,8-36].

DISCUSSION

Underlying Basic Disease

We found that most of the cases reported had AML as the basic disease followed by Acute Promyelocytic Leukaemia (APML), MDS, CML and angiogenic myeloid metaplasia. The average age of patients

was 40 years (ranging from 1.4 months to 72 years). There were in total 26 males and five females (including three children under the age 12 years). There were also four cases of post allogeneic transplantation that developed CMS during post-transplant period follow up [8].


Symptomatology

Review showed that CMS can present with a wide array of symptoms. An excellent review by Lam KY et al., of over 12000 consecutive autopsies done over the span of 20 year period showed the incidence of cardiac involvement by leukaemia of just 4% [37]. A majority of these remain undiagnosed in living individuals and discovered subsequently only on autopsy. In the symptomatic individuals, pericarditis is the frequent presentation with or without leukaemic cells in pericardial fluid [25]. Other clinical findings may include tachycardia, dyspnoea on exertion, palpitations, chest pain, arrhythmias, heart block, etc. Review showed that symptomatology of CMS in general is no more different from that of classical cardiac diseases like CAD, cardiomyopathies etc. Recognition of subtle clinical signs without frank heart failure like elevated jugular venous pressure with tall "a" wave, pedal oedema and tender hepatomegaly is crucial to warrant further imaging and invasive tests.

Diagnosis and Investigative Modalities

Diagnosis of GS requires high index of suspicion and definitive diagnosis can be very laborious and unyielding. Gold standard diagnosis is histological examination of the endomyocardial biopsy specimen [26]. Yield of biopsy samples can be increased by combining with newer diagnostic tests like MRI (SSFP/FLASH MRI) and PET scan. Superiority of PET scan is due to high efficacy in finding unifocal v/s multifocal involvement, in assessing the most 18 Fludeoxyglucose (FDG) avid site for biopsy and the treatment response after chemotherapy. Various diagnostic modalities and their utilities are summarized in [Table/Fig-3] [32,33].

A detail histopathological examination including light microscopy, immunohistochemistry and Fluorescence in Situ Hybridization (FISH) should be done on biopsy samples. The common histopathological features are highly cellular, diffuse proliferation of medium-size, polygonal to rounded, polymorph, undifferentiated malignant cells, with a high nuclear-cytoplasmic ratio, high mitotic index, scattered to diffuse areas of necrosis, and irregular production of finely reticular to coarse fibrinous tissue [34]. The cytoplasm is usually eosinophilic and granular. Positivity for CD 34, 45, 17 and HLA-DR indicates tumor mass to be of myeloid origin. Combined team effort



of radiologists, cardiologists and haematologists to diagnose and manage these challenging cases is of utmost importance.

Management Challenges

There is no consensus on the type of regimen to be used in CMS cases and haematologists worldwide have used various regimens depending on patient's tolerability and cardiac compromise. Anthracyclines are considered as backbone in the treatment therapy of AML, however the same might not be suitable in all cases due to risk of potential cardiotoxicity. Etoposide, Mitoxantrone, Cytarabine/Ara-C (EMA) based regimens have been shown to be partially efficacious in the treatment of CMS in a few case reports [35,38] Milder forms of therapy e.g., hypomethylating agents like azacytidine have also been tried which makes chemotherapy possible even in patients of older age, poor performance status and with compromised cardiac function [12,39]. A combination chemotherapy and minimal residual disease targeted therapy is needed to control the disease locally and to prevent bone marrow relapse. Newer radiotherapy protocols based on IGRT and IMRT have also broadened the treatment options where in only pathological part of heart are radiated thereby preserving the cardiac function. A total of 24 Gy in divided doses (12 fractions) was used by Yang WC et al., with successful remission achievement [12].

Author and year of publication	Age / sex	Disease /FAB Subtype	Molecular Marker/ FISH/ Cytogenetics	Cardiac chloroma at diagnosis or relapse	BM involvement	Diagnostic modality used	Pathological areas involved	Outcome
Wang Y et al., [10]	66/M	AML	NA	At diagnosis	+	PET scan 2D Echo	Ascending aorta, Pulmonary artery, pericardial effusion, B/L atrial enlargement	NA
Niu N et al., [11]	66/M	AML	NA	At diagnosis	+	PET scan 2D Echo	Both ventricles and atria, pericardium	NA
Yang WC et al., [12]	19/M	AML(Post allo-HSCT) in CR	Normal Karotype, NPM-ve)	At relapse	NA	CT scan PET scan	Soft tissue mass along the interatrial groove with an encasement of the pulmonary vein causing SVC syndrome	IMRT (24 Gy in 12 fractions) and Azacytidine following which achieved complete remission. Died soon due to sepsis.
Dorfel D et al., [13]	69/M	AML	FLT3 + ve NPM1+ve	At relapse	-	2D Echo CEMRI VPCT	Epicardial mass infiltrating right ventricular wall. Another mass at coronary sulcus	Died due to sigmoid diverticulitis and MODS
Schaffer LR et al., [14]	7/M	AML-M 5	Not mentioned	At diagnosis	+	CXR 2D Echo	Subtotal replacement of the myocardium, sinoatrial and atrioventricular (AV) nodes	Died
Kim JG et al., [15]	41/M	CML (Post MUD-HSCT) in CR	BCR-ABL 1 status not mentioned	At relapse	+	2D Echo CEMRI	Irregular shaped mass in the right atrium	Size of choloma increased after 17 months. Dose of Dasatinib was increased to 140 mg/day
Mawad R et al., [16]	42/M	Not applicable	t(8;21) 46XY	At diagnosis	-	2D Echo PET scan	Bilateral atrial walls, intra-atrial septum	Complete remission Non FDG avid residual mass
Cash T et al., [17]	24/F	AML /M5	46,del(x) Trisomy 21	At relapse	-	2D Echo CECT chest CEMRI	Mass adhering to right ventricle with RVOT obstruction pericardial effusion	Died within four months of achieving CR1 due to relapse of cholorama at the same time
Di Valentino M et al., [18]	38/F	AML	Not available	At relapse	-	2D Echo CEMRI	Infiltration of right free wall with RVOT obstruction	Complete remission
Makis W et al., [19]	42/M	AML	Not available	At diagnosis	NA	CT scan PET	Mediastinal mass with invasion of the parietal pericardium	Pericardial therapeutic drainage and HiDAC therapy- Did not achieved remission. Post two months PET showed disease progression.
Tirado CA et al., [20]	30/M	APML	+7p22, t(15;17)	At relapse	NA		Multiple sites- (left scapula, thoracic vertebra, right atrium, and supraclavicular mass	NA
Atallah A et al., [21]	56/M	AML	Not available	At diagnosis	+	2D Echo CEMRI	Focal, patchy, epicardial infiltrative process affecting mid-inferior wall	Died due to chemoresistant leukaemia

Author and year of publication	Age / sex	Disease /FAB Subtype	Molecular Marker/ FISH/ Cytogenetics	Cardiac chloroma at diagnosis or relapse	BM involvement	Diagnostic modality used	Pathological areas involved	Outcome
Tsai J et al., [22]	20/M	AML	Not available	At relapse	-	2D Echo MDCT	Well marginated, lobulated mass involving the intra-axial septum and both atria. Another mass located near the posterior aspect of tricuspid valve	Complete remission
Matkowskyj KA et al., [23]	59/M	T-MDS – RAEB type (treated 3 years back for Anaplastic oligodendroglioma)	Abnormal mosaic male karyotype (46,XY and 45,XY,-7)	At diagnosis	-	2D Echo	Left ventricle, septum, and right ventricle	Died before treatment could be started
Mignano JE [24]	20/M	AML, FAB M-2	Not mentioned	Relapse	-	CT	Right and left atria, intra-atrial septum, and apparent extension into the right pulmonary veins	Resolution
Tsai MH [25]	1.4 mon/F	AML	51,XX,add(1)(p32),del(4)(q31),+del(6)(q23),add(7)(q11.2),+8,+8,der(11)t(7;11)(q11.2;q23),+13,add(16)(q24),+19.	At diagnosis	+	CXR, ECG	Pericardium invasion	Died secondary to sepsis
Rigamonti F et al., [27]	52/M	AML	47,X,add(Y)(q12)	At diagnosis	-	Cardiac Ultrasound	Infiltration of the infero-lateral cardiac wall, right auricle and aortic arch	3+7 Regimen/ Patient died despite achieving remission due to fungal sepsis
Kozelj M [26]	52/M	AML	t(8;21),	At diagnosis	+	2D Echo, TTE, MRI	Right atrium, right pulmonary hilum	Remission
Antic D et al., [28]	37/M	AML	NA	At diagnosis	-	CT Chest	Large mass in right atrium extending into SVC causing SVC syndrome	Surgical excision of the mass followed by 3+7 regimen
Diab M et al., [29]	27/M	AML	NA	At diagnosis	-	CT Chest	Leukemia cutis, Pericardial effusion, mediastinal mass	Pericardiocentesis, Cytrabine, Daunorubicin and etoposide based regimen. Refractory to chemotherapy due to which patient soon.
Nasilowska AB et al., [30]	34/M	APML in CR (Post allo-HSCT)	t(15;17)	At Relapse	-	CT Chest	Multiple sites of EM relapse- pleura, heart and pericardium	During salvage chemotherapy, patient died of cardiogenic shock
Mateen FJ et al., [1]	64/W	MDS, RAEB	Not mentioned	At diagnosis	+	CXR, 2D echo	Myocardium infiltration	Died
Antón E et al., [31]	45/F	Not applicable	+4, -9, -17, add(17)(p13), -18, +21, +22	At diagnosis	-	CXR, ECG	Myocardial infiltration	Died due to multi organ failure
Kara IO et al., [8]	28/M	AML, FAB-M2 in CR 2 (Post allo-HSCT)	Not mentioned	At relapse	+	2D Echo	Nasopharynx nodular masses, right atrium, right ventricle	Etoposide,mitoxantrone, ara-C (EMA) chemotherapy regimen-achieved CR
Marcos AP et al., [32]	39/M	AML, FAB-M1	Not mentioned	At relapse	-	TEE	Right atrium	Complete remission
Markaryus AN et al., [33]	34/M	AML, FAB-M1	Not mentioned	At relapse	+	TTE, 2D echo, TEE, MRI	All chambers	Died
Erdol C et al., [34]	48/M	AML, FAB-M4	Not mentioned	At diagnosis	+	TTE, TEE, Echocardiography	Right atrium	Complete regression post chemotherapy
Tillawi IS et al., [35]	72/M	Angiogenic myeloid metaplasia			NA		Encasement of the heart and great vessels	
Jankovic M et al., [9]	12/F	APML	Not done	At relapse	-	CXR, Echocardiography	Right-sided intracardiac mass, pericardial effusion	Died due to progressive heart failure
Foucar K et al., [36]	22/M	Not applicable	Not done	At diagnosis	-	CXR, Echocardiography	Encasement of the SVC, PA, and aorta	Died due to ventricular arrhythmia

[Table/Fig-2]: Review of published cases of cardiac chloromas in literature from 1960 till July 2016, their baseline disease characters, molecular characteristics, diagnostic modality used, areas of cardiac tissue involved with outcome [1,8-36].

CONCLUSION

Despite various chemotherapeutic and radiation based treatment protocols, CMS carries a dismal prognosis. High mortality and morbidity is associated with this rare disease. Major limitation is

that anthracycline based regimens cannot be used due to their own cardiotoxicity. Further research is warranted to identify newer molecular markers for diagnosis, standardize diagnostic tests and formulate efficacious chemotherapy and radiation protocols.

ECG, Chest X Ray	Non-specific test, cheap, beneficial to give quick idea about arrhythmia, effusion and cardiac borders and mediastinal involvement.
2D - Echocardiography and TEE	Fast and dynamic, TEE can be used to take biopsy.
CT Scan	Quick, easily available. To analyze intra-cardiac invasion and involvement of pulmonary vasculature. Volume perfusion scan (VPCT) to assess viable myocardium.
MRI	Steady state free precession imaging (SSFP MRI) and fast low angle short magnetic resonance imaging (FLASH MRI) can acquire dynamic images synchronised with ECG.
PET Scan [40]	Best to assess the site for biopsy and the treatment response.

[Table/Fig-3]: Diagnostic modalities and their efficacy in diagnosing.

REFERENCES

- Mateen FJ, Harding SR, Saxena A. Extensive myocardial infiltration by hemopoietic precursors in a patient with myelodysplastic syndrome. *BMC Blood Disord.* 2006;6:4.
- Sahu KK, Yanamandra U, Malhotra P. Orbital myeloid sarcoma: Rare presentation of AML. *Orbit.* 2016;35(3):157–58.
- Sahu KK, Malhotra P. Re: "Granulocytic sarcoma of the orbit presenting as a fulminant orbitopathy in an adult with acute myeloid leukaemia". *Ophthal Plast Reconstr Surg.* 2015;31(5):421.
- Sahu KK, Jain A, Yanamandra U, Varma SC, Malhotra P. Myeloid sarcoma of vulva: a short update. *Indian J Haematol Blood Transfus an Off J Indian Soc Haematol Blood Transfus.* 2016;32(Suppl 1):69–71.
- Jain A, Sahu KK, Sharma S, Rajwanshi A, Suri V, Malhotra P. Shoulder myeloid sarcoma: an initial presentation of cml blast crisis. *Indian J Haematol blood Transfus an Off J Indian Soc Haematol Blood Transfus.* 2016;32(Suppl 1):361–63.
- Sahu KK, Sanamandra P, Jeyaraman P, Kumar G, Prakash G, Kumar N, et al. Unusual cause of cord compression-A pressing issue for neurosurgeons. *World Neurosurg.* 2016;92:565–67.
- Sahu KK, Tyagi R, Law AD, Khadwal A, Prakash G, Rajwanshi A, et al. Myeloid sarcoma: an unusual case of mediastinal mass and malignant pleural effusion with review of literature. *Indian J Haematol blood Transfus an Off J Indian Soc Haematol Blood Transfus.* 2015;31(4):466–71.
- Kara IO, Sahin B, Paydas S, Kara B. Granulocytic sarcoma of the heart: extramedullary relapse of acute myeloblastic leukaemia after allogeneic stem cell transplantation successfully treated by chemotherapy alone. *Leuk Lymphoma.* 2005;46(7):1081–84.
- Jankovic M, Bonacina E, Masera G, Uderzo C, Galli MA, Ottaviani V, et al. Cardiac relapses in myeloid leukaemia: case report and review of the literature. *Pediatr Haematol Oncol.* 1987;4(3):237–45.
- Wang Y, Wu W, Zhong D, Han X, Fang L. Multimodality imaging and histopathology of cardiac myeloid sarcoma. *Eur Heart J Cardiovasc Imaging.* 2016;17(11):1316.
- Niu N, Cui R, Li F. FDG PET/CT findings of intracardiac myeloid sarcoma. *Clin Nucl Med.* 2016;41(3):235–36.
- Yang WC, Yao M, Chen YH, Kuo SH. Complete response of myeloid sarcoma with cardiac involvement to radiotherapy. *J Thorac Dis.* 2016;8(6):1323–28.
- Dorfel D, Hantschel M, Federmann B, Haen S, Fend F, Muller II, et al. Cardiac myeloid sarcoma: multimodality radiologic imaging features and pathologic correlation. *Am J Med.* 2016;129(8):e117–20.
- Schaffer LR, Caltharp SA, Milla SS, Kogon BF, Cundiff CA, Dalal A, et al. Rare presentation of four primary pediatric cardiac tumors. *Cardiovasc Pathol.* 2016;25(1):72–77.
- Kim JG, Moon D, Yi JE, Youn HJ, Kim DW, Park GS et al. Recurrent cardiac chloroma presenting as acute chest pain. *QJM.* 2014;107(5):381–82.
- Mawad R, Wu D, Abkowitz JL, Walter RB. Myeloid sarcoma of the heart. *Leuk lymphoma.* 2012;53:2511–14.
- Cash T, Becton D, Mian A. Cardiac myeloid sarcoma: a case report and review of literature. *J Pediatr Haematol Oncol.* 2011;33(7):e330–32.
- Di Valentino M, Menafoglio A, Wyrtenbach R, Galliono A. An unusual cause of recurrent syncope. *J Am Coll Cardiol.* 2011;58(16):1728.
- Makis W, Hickeson M, Derbeykan V. Myeloid sarcoma presenting as an anterior mediastinal mass invading the pericardium: Serial imaging with F-18 FDG PET/CT. *Clin Nucl Med.* 2010;35(9):706–09.
- Tirado CA, Chen W, Vadez F, Krandikar N, Arbini A, Acevedo I, et al. Unusual presentation of myeloid sarcoma in a case of acute pyelomyelocytic leukaemia with a cryptic PML-RARA rearrangement involving multiple sites including the atrium. *Cancer Gnet Cytogenet.* 2010;200(1):47–53.
- Atallah A, Cheong BY, Bernicker E, Wilson JM. Cardiac chloroma: novel presentation and subsequent diagnosis with cardiac magnetic resonance imaging. *Tex Heart Inst J.* 2010;37(2):242–43.
- Tsai J, Lee EY. MDCT imaging findings of extramedullary granulocytic sarcoma of the heart. *J Thorac Imaging.* 2010;25(1):W14–16.
- Matkowskyj KA, Wiseman WR, Robin JC, Norvell JP, Puthumana J, Nelson B, et al. Therapy related myelodysplastic syndrome presenting as fulminant heart failure secondary to myeloid sarcoma. *J Haematop.* 2010;3(1):41–46.
- Mignano JE, Chan MD, Rosenwald IB, Kimmelstiel CD, Wolfe LC. Intracardiac chloroma. *J Pediatr Haematol Oncol.* 2009;31(12):977–79.
- Tsai MH, Yang CP, Chung HT, Shih LY. Acute myeloid leukaemia in a young girl presenting with mediastinal granulocytic sarcoma invading pericardium and causing superior vena cava syndrome. *J Pediatr Haematol Oncol.* 2009;31(12):980–82.
- Kozelj M, Zorman D, Mrevlje B, Cernelc P, Zver S. Cardiac granulocytic sarcoma diagnosed by intracardiac echocardiography guided biopsy. *Int J Haematol.* 2008;88(1):101–03.
- Rigamonti F, Beris P, Sanchez Pareja A, Meyer P, Ashrafpoor G, et al. Atypical presentation of acute myeloid leukaemia: cardiac myeloid sarcoma. *Int J Haematol.* 2009;89(5):693–98.
- Antic D, Vuckovic M, Elezovic I. Right atrial myeloid sarcoma causing superior vena cava syndrome. *Br J haematol.* 2008;141(2):134.
- Diab M, Coloe J, Bechtel M. Extramedullary granulocytic sarcoma of the skin, mediastinum and pericardium. *Int J Dermatol.* 2008;47(3):256–58.
- Nasilowska -Adamska B, Majewski M, Seferynska I, Szczepinski A, Tomaszewska A, ProchrocecSobeiszek M, et al. Predictive value of RT-PCR PML-RARA transcript monitoring for extramedullary relapse of acute promyelocytic leulemia in the pleura, heart and pericardium after allogeneic SCT. *Ann Transplant.* 2007;12(3):33–38.
- Anton E. Cardiovascular involvement in adult granulocytic sarcoma. *Am J Haematol.* 2006;81(5):382–83.
- Marcos-Alberca P1, Ibañez B, Rey M, Román A, Rábago R, Orejas M, et al. Cardiac granulocytic sarcoma (chloroma): in vivo diagnosis with transesophageal echocardiography. *J Am Soc Echocardiogr.* 2004;17(9):1000–02.
- Makaryus AN, Tung F, Liu W, Mangion J, Kort S. Extensive neoplastic cardiac infiltration in a patient with acute myelogenous leukaemia: role of echocardiography. *Echocardiography.* 2003;20(6):539–44.
- Erdol C, Ovali E, Baykan M. Granulocytic sarcoma presenting as a right atrial mass. *Acta Cardiol.* 2003;58(2):155–58.
- Tillawi IS, Variakojis ID. Refractory right ventricular failure due to granulocytic sarcoma. *Arch Pathol Lab Med.* 1990;114(9):983–85.
- Foucar K, Foucar E, Willman C, Horvath A, Gerety RL. Nonleukaemic granulocytic sarcoma of the heart: A report of a fatal case. *Am J Haematol.* 1987;25(3):325–32.
- Lam KY, Dickens P, Chan AC. Tumors of the heart. A 20-year experience with a review of 12,485 consecutive autopsies. *Arch Pathol Lab Med.* 1993;117(10):1027–31.
- Tabbane C, Chadly A, Corcos V, Heldt. A case of chloroma with cardiac metastasis in an 8-year-old child. *Tunis Med.* 1960;38:153–64.
- Sahu KK, Dhbar DP, Malhotra P. Isolated myeloid sarcoma. *Orbit.* 2016;35(6):351.
- Sahu KK, Gautam A, Ailawadhi S. Re: FDG PET/CT findings of intracardiac myeloid sarcoma. *Clin Nucl Med.* 2017 Mar;42(3):242–45.

PARTICULARS OF CONTRIBUTORS:

- Junior Resident, Department of Internal Medicine, UCMS, New Delhi, India.
- Senior Resident, Department of Internal Medicine, Shahid Beheshti University of Medical Science, Tehran, Iran.
- Senior Resident, Department of Internal Medicine, PGIMER, Chandigarh, India.
- Junior Resident, Department of Internal Medicine, PGIMER, Chandigarh, India.
- Senior Associate Consultant, Department of Internal Medicine, Mayo Clinic, Jacksonville, Florida.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Kamal Kant Sahu,
Senior Resident, Department of Internal Medicine, Nehru Hospital, PGIMER, Sector 12, Pin-160012, Chandigarh, India.
E-mail: everlastingkamal@yahoo.com

Date of Submission: **Aug 02, 2016**
Date of Peer Review: **Aug 24, 2016**
Date of Acceptance: **Jan 23, 2017**
Date of Publishing: **Mar 01, 2017**

FINANCIAL OR OTHER COMPETING INTERESTS: None.